




PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07D 215/14, A61K 31/395, C07D 277/64, 215/18, 213/30, 241/42, 235/12	A1	(11) International Publication Number: WO 97/12867 (43) International Publication Date: 10 April 1997 (10.04.97)
(21) International Application Number: PCT/US96/15345 (22) International Filing Date: 25 September 1996 (25.09.96) (30) Priority Data: 60/004,706 3 October 1995 (03.10.95) US 08/703,441 17 September 1996 (17.09.96) US (71) Applicant: ABBOTT LABORATORIES [US/US]; CHAD 0377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-3500 (US). (72) Inventors: BROOKS, Clint, D., W.; 928 Quaker Hill Lane, Libertyville, IL 60048 (US). BHATIA, Pramila; 1602 Mulberry Drive, Libertyville, IL 60048 (US). KOLASA, Theodozyj; 118 Walden Lane, Lake Villa, IL 60046 (US). STEWART, Andrew, O.; 1289 Thornbury Lane, Liber- tyville, IL 60048 (US). GUNN, David, E.; 40 Wood Street, Hamden, CT 06517 (US). CRAIG, Richard, A.; 3212 Fox Drive, Racine, WI 53404 (US). (74) Agents: ANAND, Mona et al.; Abbott Laboratories, CHAD 0377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-3500 (US).	(81) Designated States: AU, CA, IL, JP, KR, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>	
(54) Title: SYMMETRICAL bis-HETEROARYLMETHOXYPHENYLALKYL CARBOXYLATES AS INHIBITORS OF LEUKOTRIENE BIOSYNTHESIS <div style="text-align: center;">  </div> (57) Abstract <p>Compounds having formula (I), wherein W is the same at each occurrence and is selected from optionally substituted quinolyl, optionally substituted benzothiazolyl, optionally substituted benzoxazolyl, optionally substituted benzimidazolyl, optionally substituted quinoxalyl, optionally substituted pyridyl, optionally substituted pyrimidyl, and optionally substituted thiazolyl; R¹ and R² are independently selected from hydrogen, alkyl, haloalkyl, alkoxy, halogen; R³ is a valence bond or is selected from hydrogen and alkyl; X is a valence bond or is selected from alkylene, alkenylene, and alkynylene; and Z is selected from (a) COM, (b) CH=N-O-A-COM, (c) CH₂-O-N-A-COM wherein A is selected from alkylene and cycloalkylene, and M is selected from (a) a pharmaceutically acceptable metabolically cleavable group, (b) -OR⁶, (c) -NR⁷R⁸, (d) -NR⁶SO₂R⁹, (e) -NH-Tetrazolyl, and (f) glycyl inhibit leukotriene biosynthesis and are useful in the treatment of allergic and inflammatory disease states. Also disclosed are leukotriene biosynthesis inhibiting compositions and a method of inhibiting leukotriene biosynthesis.</p>		

Best Available Copy

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**SYMMETRICAL *bis*-HETEROARYLMETHOXY-
PHENYLALKYL CARBOXYLATES AS INHIBITORS
OF LEUKOTRIENE BIOSYNTHESIS**

Cross Reference To Related Applications

This application claims benefit of U.S. Provisional Application No. 60/004,706, filed October 3, 1995.

5 Technical Field

This invention relates to compounds having activity to inhibit leukotriene biosynthesis, to pharmaceutical compositions comprising these compounds, and to a medical method of treatment. More particularly, this invention concerns a class of symmetrical *bis*-heteroarylmethoxyphenylalkyl carboxylate compounds which inhibit
10 leukotriene biosynthesis, to pharmaceutical compositions comprising these compounds and to a method of inhibiting leukotriene biosynthesis.

Background Of The Invention

The leukotrienes are extremely potent substances which produce a wide
15 variety of biological effects, often even when present only in nanomolar to picomolar concentrations. Leukotrienes are important pathological mediators in a variety of diseases. Alterations in leukotriene metabolism have been demonstrated in a number of disease states including asthma, allergic rhinitis, rheumatoid arthritis and gout, psoriasis, adult respiratory distress syndrome, inflammatory bowel disease,
20 endotoxin shock syndrome, atherosclerosis, ischemia induced myocardial injury, and central nervous system pathology resulting from the formation of leukotrienes following stroke or subarachnoid hemorrhage.

Compounds which prevent leukotriene biosynthesis are thus useful in the treatment of disease states such as those listed above in which the leukotrienes play an
25 important pathophysiological role.

United States Patent Number 5,358,955 (October 25, 1994) discloses aryl and heteroarylmethoxyphenyl compounds which inhibit leukotriene biosynthesis.

United States Patent Number 4,970,215 (November 13, 1990) discloses quinolylmethoxyphenyl acetic acid derivatives which inhibit leukotriene biosynthesis.

30 United States Patent Number 5,512,581 discloses iminoxycarboxylate derivatives which inhibit leukotriene biosynthesis. United States Patent Number 5,326,883 (July 5, 1994) discloses oxime ether derivatives having lipoxigenase inhibitory activity.


European Patent Application Number 349 062 (January 3, 1990) discloses quinolylmethoxyphenyl alkanolic acid derivatives which inhibit leukotriene biosynthesis.

PCT Application Number WO 94/27968 (December 8, 1994) discloses
5 quinoline derivatives as leukotriene antagonists.

Prasit, *et al.*, *Bioorganic and Medicinal Chemistry Letters*, 1 (11), 645 (1991) describe ((4-(4-chlorophenyl)-1-(4-(2-quinolylmethoxy)phenyl)butyl)thio)acetic acid as an orally active leukotriene biosynthesis inhibitor, and Musser and Kraft, *Journal of Medicinal Chemistry*, 35 (14), 1, (1992) review quinoline containing leukotriene
10 biosynthesis inhibitors.

Summary Of The Invention

In its principal embodiment, the present invention provides a class of novel symmetrical *bis*-heteroarylmethoxyphenylalkyl carboxylate compounds and their
15 derivatives and pharmaceutically acceptable salts. The compounds have the formula:



wherein **W** is the same at each occurrence and is selected from the group consisting of
20 (a) quinolyl; (b) quinolyl substituted with a substituent selected from the group consisting of (b-1) halogen, (b-2) alkyl of one to six carbon atoms, (b-3) phenyl, (b-4) phenyl substituted with a substituent selected from the group consisting of (b-4-a) halogen, (b-4b) alkyl of one to six carbon atoms, (b-4-c) haloalkyl of one to six carbon atoms, and (b-4-d) alkoxy of one to six carbon atoms, (b-5) pyridyl, and (b-6)
25 pyridyl substituted with a substituent selected from the group consisting of (b-6-a) halogen, (b-6-b) alkyl of one to six carbon atoms, and (b-6-c) alkoxy of one to six carbon atoms; (c) benzothiazolyl; (d) benzothiazolyl substituted with a substituent selected from the group consisting of (d-1) halogen, (d-2) alkyl of one to six carbon atoms, (d-3) phenyl, (d-4) phenyl substituted with a substituent selected from the
30 group consisting of (d-4-a) halogen, (d-4-b) alkyl of one to six carbon atoms, (d-4-c) haloalkyl of one to six carbon atoms, and (d-4-d) alkoxy of one to six carbon atoms, (d-5) pyridyl, and (d-6) pyridyl substituted with a substituent selected from the group consisting of (d-6-a) halogen, (d-6-b) alkyl of one to six carbon atoms, and (d-6-c)

alkoxy of one to six carbon atoms; (e) benzoxazolyl; (f) benzoxazolyl substituted with a substituent selected from the group consisting of (f-1) halogen, (f-2) alkyl of one to six carbon atoms, (f-3) phenyl, (f-4) phenyl substituted with a substituent selected from the group consisting of (f-4-a) halogen, (f-4-b) alkyl of one to six carbon atoms, (f-4-c) haloalkyl of one to six carbon atoms, and (f-4-d) alkoxy of one to six carbon atoms, (f-5) pyridyl, and (f-6) pyridyl substituted with a substituent selected from the group consisting of (f-6-a) halogen, (f-6-b) alkyl of one to six carbon atoms, and (f-6-c) alkoxy of one to six carbon atoms; (g) benzimidazolyl; (h) benzimidazolyl substituted with a substituent selected from the group consisting of (h-1) halogen, (h-2) alkyl of one to six carbon atoms, (h-3) phenyl, (h-4) phenyl substituted with a substituent selected from the group consisting of (h-4-a) halogen, (h-4-b) alkyl of one to six carbon atoms, (h-4-c) haloalkyl of one to six carbon atoms, and (h-4-d) alkoxy of one to six carbon atoms, (h-5) pyridyl, and (h-6) pyridyl substituted with a substituent selected from the group consisting of (h-6-a) halogen, (h-6-b) alkyl of one to six carbon atoms, and (h-6-c) alkoxy of one to six carbon atoms; (i) quinoxalyl; (j) quinoxalyl substituted with a substituent selected from the group consisting of (j-1) halogen, (j-2) alkyl of one to six carbon atoms, (j-3) phenyl, (j-4) phenyl substituted with a substituent selected from the group consisting of (j-4-a) halogen, (j-4-b) alkyl of one to six carbon atoms, (j-4-c) haloalkyl of one to six carbon atoms, and (j-4-d) alkoxy of one to six carbon atoms, (j-5) pyridyl, and (j-6) pyridyl substituted with a substituent selected from the group consisting of (j-6-a) halogen, (j-6-b) alkyl of one to six carbon atoms, and (j-6-c) alkoxy of one to six carbon atoms; (k) pyridyl; (l) pyridyl substituted with a substituent selected from the group consisting of (l-1) phenyl, (l-2) phenyl substituted with a substituent selected from the group consisting of (l-2-a) halogen, (l-2-b) alkyl of one to six carbon atoms, (l-2-c) haloalkyl of one to six carbon atoms, and (l-2-d) alkoxy of one to six carbon atoms, (l-3) pyridyl, and (l-4) pyridyl substituted with a substituent selected from the group consisting of (l-4-a) halogen, (l-4-b) alkyl of one to six carbon atoms, and (l-4-c) alkoxy of one to six carbon atoms; (m) pyrimidyl; (n) pyrimidyl substituted with a substituent selected from the group consisting of (n-1) phenyl, (n-2) phenyl substituted with a substituent selected from the group consisting of (n-2-a) halogen, (n-2-b) alkyl of one to six carbon atoms, (n-2-c) haloalkyl of one to six carbon atoms, and (n-2-d) alkoxy of one to six carbon atoms, (n-3) pyridyl, and (n-4) pyridyl substituted with a substituent selected from the group consisting of (n-4-a) halogen, (n-4-b) alkyl of one to six carbon atoms, and (n-4-c) alkoxy of one to six carbon atoms; (o) thiazolyl, and (p) thiazolyl substituted with a substituent selected from the group consisting of (p-1) phenyl, (p-2) phenyl substituted with a substituent selected from the group consisting

of (p-2-a) halogen, (p-2-b) alkyl of one to six carbon atoms, (p-2-c) haloalkyl of one to six carbon atoms, and (p-2-d) alkoxy of one to six carbon atoms, (p-3) pyridyl, and (p-4) pyridyl substituted with a substituent selected from the group consisting of (p-4-a) halogen, (p-4-b) alkyl of one to six carbon atoms, and (p-4-c) alkoxy of one to six carbon atoms.

R^1 and R^2 are independently selected from the group consisting of (a) hydrogen, (b) alkyl of one to six carbon atoms, (c) haloalkyl of one to six carbon atoms, (d) alkoxy of one to six carbon atoms, and (e) halogen.

R^3 is selected from the group consisting of (a) hydrogen and (b) alkyl of one to six carbon atoms.

X is absent or is selected from the group consisting of (a) alkylene of one to six carbon atoms, (b) alkenylene of one to six carbon atoms, and (c) alkynylene of one to six carbon atoms.

Z is selected from the group consisting of (a) COM, (b) CH=N-O-A-COM, (c) CH₂-O-N=A-COM, and (d) OR³ where R^3 is hydrogen or alkyl of one to six carbon atoms.

A is selected from the group consisting of (a) alkylene of one to six carbon atoms, and (b) cycloalkylene of three to eight carbon atoms.

M is selected from the group consisting of (a) a pharmaceutically acceptable metabolically cleavable group, (b) -OR³ where R^3 is selected from the group consisting of hydrogen and alkyl of one to six carbon atoms, and (c) -NR⁷R⁸ where R^7 and R^8 are independently selected from the group consisting of hydrogen, alkyl of one to six carbon atoms, hydroxy, and alkoxy of one to six carbon atoms, or R^7 and R^8 taken together define a five- to eight-membered ring, with the proviso that R^7 and R^8 may not simultaneously be hydroxyl, (d) -NR³SO₂R⁹ wherein R^3 is as defined above and R^9 is alkyl of one to six carbon atoms, (e) -NH-tetrazolyl, and (f) glycyl.

The present invention also provides pharmaceutical compositions which comprise a therapeutically effective amount of compound as defined above in combination with a pharmaceutically acceptable carrier.

The invention further relates to a method of inhibiting leukotriene biosynthesis in a host mammal in need of such treatment comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound as defined above.

Detailed Description

As used throughout this specification and the appended claims, the following terms have the meanings specified.

The term alkyl refers to a monovalent group derived from a straight or branched chain saturated hydrocarbon by the removal of a single hydrogen atom. Alkyl groups are exemplified by methyl, ethyl, n- and iso-propyl, *n*-, *sec*-, *iso*- and *tert*-butyl, and the like.

- 5 The terms alkoxy and alkoxyl denote an alkyl group, as defined above, attached to the parent molecular moiety through an oxygen atom. Representative alkoxy groups include methoxy, ethoxy, propoxy, butoxy, and the like.

- 10 The term alkenyl as used herein refers to monovalent straight or branched chain groups of 2 to 6 carbon atoms containing a carbon-carbon double bond, derived from an alkene by the removal of one hydrogen atom and include, but are not limited to groups such as ethenyl, 1 -propenyl, 2-propenyl, 2-methyl- 1 -propenyl, 1 -butenyl, 2-butenyl and the like.

- 15 The term alkylene denotes a divalent group derived from a straight or branched chain saturated hydrocarbon by the removal of two hydrogen atoms, for example -CH₂-, -CH₂CH₂-, -CH(CH₃)CH₂- and the like.

The term alkenylene denotes a divalent group derived from a straight or branched chain hydrocarbon containing at least one carbon-carbon double bond. Examples of alkenylene include -CH=CH-, -CH₂CH=CH-, -C(CH₃)=CH-, -CH₂CH=CHCH₂-, and the like.

- 20 The term alkynylene refers to a divalent group derived by the removal of two hydrogen atoms from a straight or branched chain acyclic hydrocarbon group containing at least one carbon-carbon triple bond. Examples of alkynylene include —CH≡CH—, —CH≡CH-CH₂—, —CH≡CH-CH(CH₃)—, and the like.

- 25 The term aryl as used herein refers to a monovalent carbocyclic group containing one or more fused or non-fused phenyl rings and includes, for example, phenyl, 1- or 2-naphthyl, 1,2-dihydronaphthyl, 1,2,3,4-tetrahydronaphthyl, and the like.

- 30 The term cycloalkyl as used herein refers to a monovalent saturated cyclic hydrocarbon group. Representative cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, bicyclo[2.2.1]heptane and the like.

Cycloalkylene denotes a divalent radical derived from a cycloalkane by the removal of two hydrogen atoms.

- 35 The term haloalkyl denotes an alkyl group, as defined above, having one, two, or three halogen atoms attached thereto and is exemplified by such groups as chloromethyl, bromoethyl, trifluoromethyl, and the like.

As used throughout this specification and the appended claims, the term "metabolically cleavable group" denotes a moiety which is readily cleaved *in vivo* from the compound bearing it, which compound after cleavage remains or becomes pharmacologically active. Metabolically cleavable groups form a class of groups reactive with the carboxyl group of the compounds of this invention (where M is -OH) well known to practitioners of the art. They include, but are not limited to such groups as alkanoyl (such as acetyl, propionyl, butyryl, and the like), unsubstituted and substituted aroyl (such as benzoyl and substituted benzoyl), alkoxycarbonyl (such as ethoxycarbonyl), trialkylsilyl (such as trimethyl- and triethylsilyl), monoesters formed with dicarboxylic acids (such as succinyl), and the like. Because of the ease with which the metabolically cleavable groups of the compounds of this invention are cleaved *in vivo*, the compounds bearing such groups act as pro-drugs of other leukotriene biosynthesis inhibitors. The compounds bearing the metabolically cleavable groups have the advantage that they may exhibit improved bioavailability as a result of enhanced solubility and/or rate of absorption conferred upon the parent compound by virtue of the presence of the metabolically cleavable group.

In those instances where M = OH, the compounds of the present invention are capable of forming base addition salts. In such instances, the term "pharmaceutically acceptable salts" refers to the relatively nontoxic inorganic and organic base addition salts of compounds of the present invention. These salts can be prepared *in situ* during the final isolation and purification of the compounds, or by separately reacting the purified carboxyl compound with a suitable base such as the hydroxide, carbonate, or bicarbonate of a pharmaceutically acceptable metal cation or with ammonia, or an organic primary, secondary, or tertiary amine of sufficient basicity to form a salt with the carboxyl functional group of the compounds of this invention.

Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Representative organic amines useful for the formation of base addition salts include ethylamine, diethylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine, and the like. (See, for example, S. M. Berge, *et al.*, *J. Pharmaceutical Sciences*, 1977, 66: 1-19, which is incorporated herein by reference).

Similarly, in those instances where the compounds of the present invention possess a heterocyclic ring moiety containing a basic nitrogen atom, the compounds are capable of forming acid addition salts. In such cases, the term "pharmaceutically acceptable salts" also refers to the nontoxic inorganic and organic acid addition salts of compounds of the present invention. These salts can be prepared *in situ* during the final isolation and purification of the compounds, or by separately reacting the

purified compound in its free-base form with a suitable inorganic or organic acid and isolating the salt thus formed. Representative acid addition salts include acetate, adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphersulfonate, citrate, cyclopentanepropionate, digluconate, 5 dodecylsulfate, ethanesulfonate, fumarate, glucoheptonate, glycerophosphate, hemisulfate, heptonate, hexanoate, hydrobromide, hydrochloride, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pantoate, pectinate, persulfate, 3-phenylpropionate, phosphate, 10 picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, toluenesulfonate, undecanoate, valerate salts, and the like. (See, for example, S. M. Berge, *et al.*, *J. Pharmaceutical Sciences*, **1977**, 66: 1-19), which is incorporated herein by reference). Said pharmaceutically acceptable acid and base addition salts are also contemplated as falling within the scope of the present invention.

15 Asymmetric centers may exist in the compounds of the present invention. The present invention contemplates the various stereoisomers and mixtures thereof. Individual stereoisomers of compounds of the present invention are made by synthesis from starting materials containing the chiral centers or by preparation of mixtures of enantiomeric products followed by separation as, for example, by 20 conversion to a mixture of diastereomers followed by separation by recrystallization or chromatographic techniques, or by direct separation of the optical enantiomers on chiral chromatographic columns. Starting compounds of particular stereochemistry are either commercially available or are made by the methods detailed below and resolved by techniques well known in the organic chemical arts.

25 Examples of compounds contemplated as falling within the scope of the present invention include, but are not limited to:

4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid,
4,4-*bis*(4-(2-quinolylmethyl)phenyl)pentanoic acid methyl ester,
4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid sodium salt,
30 4,4-*bis*(4-(2-quinolylmethoxy)phenyl) pentanoic acid magnesium salt,
4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentan-1-ol,
[4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pent-1-yl]iminoxyacetic acid,
[4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pent-1-yl]-2-iminoxypropionic acid,
[4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pent-1-yl]-2-iminoxypropionic acid,
35 methyl ester,
[4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pent-1-yl]oximinoacetic acid,
4,4-*bis*(4-(2-benzothiazoylmethoxy)phenyl)pentanoic acid,

- 4,4-*bis*(4-(2-benzothiazolylmethoxy)phenyl)pentanoic acid sodium salt,
 4,4-*bis*(4-(7-chloro-2-quinolylmethoxy)phenyl)pentanoic acid,
 4,4-*bis*(4-(6-fluoro-2-quinolylmethoxy)phenyl)pentanoic acid sodium salt,
 4,4-*bis*(4-(6-fluoro-2-quinolylmethoxy)phenyl)pentanoic acid,
 5 2,2-*bis*(4-(2-quinolylmethoxy)phenyl)propionic acid,
 3,3-*bis*(4-(2-quinolylmethoxy)phenyl)butanoic acid,
 5,5-*bis*(4-(2-quinolylmethoxy)phenyl)hexanoic acid,
 5,5-*bis*(4-(2-quinolylmethoxy)phenyl)hexanoic acid sodium salt,
 4,4-*bis*(4-(2-pyridylmethoxy)phenyl)pentanoic acid sodium salt,
 10 4,4-*bis*(4-(2-pyridylmethoxy)phenyl)pentanoic acid,
 4,4-*bis*(4-(2-quinoxylmethoxy)phenyl)pentanoic acid sodium salt,
 4,4-*bis*(4-(1-methyl-2-benzimidazolylmethoxy)phenyl)pentanoic acid,
 4,4-*bis*(4-(1-methyl-2-benzimidazolylmethoxy)phenyl)pentanoic acid sodium
 salt,
 15 4,4-*bis*(3-chloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid,
 4,4-*bis*(3-chloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid methyl ester,
 4,4-*bis*(3-chloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid sodium salt,
 [4,4-*bis*(3-chloro-4-(2-quinolylmethoxy)phenyl)pent-1-yl]-2-
 iminoxypionic acid sodium salt,
 20 4,4-*bis*(3-chloro-4-(2-quinolylmethoxy)phenyl)pentan-1-ol,
 [4,4-*bis*(3-chloro-4-(2-quinolylmethoxy)phenyl)pent-1-yl]-2-
 iminoxypionic acid,
 4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid *N,N*-
 diethylhydroxylamine ester,
 25 2,2-*bis*(4-(2-quinolylmethoxy)phenyl)butyric acid,
 1,1-*bis*(4-(2-quinolylmethoxy)phenyl)ethanol,
 2,2-dimethyl-3,3-*bis*(4-(2-quinolylmethoxy)phenyl)propionic acid sodium
 salt,
 2,2-dimethyl-3,3-*bis*(4-(2-quinolylmethoxy)phenyl)propionic acid methyl
 30 ester,
 2,2-dimethyl-3,3-*bis*(4-(2-quinolylmethoxy)phenyl)propionic acid,
 [2,2-dimethyl-3,3-*bis*(4-(2-quinolylmethoxy)phenyl)prop-1-yl]oximinoacetic
 acid sodium salt,
 2,2-dimethyl-3,3-*bis*(4-(2-quinolylmethoxy)phenyl)propan-1-ol,
 35 [2,2-dimethyl-3,3-*bis*(4-(2-quinolylmethoxy)phenyl)prop-1-yl]oximinoacetic
 acid,

2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)-1-propyliminoxyacetic acid,

2,2-bis(4-(2-quinolylmethoxy)phenyl)acetic acid,

2,2-bis(4-(6-fluoro-2-quinolylmethoxy)phenyl)acetic acid,

5 2,2-bis(4-(2-quinolylmethoxy)phenyl)eth-1-yloximinoacetic acid,

2,2-bis(4-(2-quinolylmethoxy)phenyl)acetic acid methyl ester,

2,2-bis(4-(2-quinolylmethoxy)phenyl)eth-1-yloximinoacetic acid sodium salt,

3,3-bis(4-(2-quinolylmethoxy)phenyl)propionic acid,

3,3-bis(4-(2-quinolylmethoxy)phenyl)propionic acid sodium salt,

10 4,4-bis(4-(2-quinolylmethoxy)phenyl)acetic acid-*N*-carboxymethyl amide,

[3,3-bis(2-quinolylmethoxyphenyl)but-1-yl]-2-iminoxypropionic acid,

3,3-bis(4-(2-quinolylmethoxy)phenyl)butan-1-ol,

4,4-bis(4-(2-quinolylmethoxy)phenyl)-4-hydroxy-2-butynoic acid,

4,4-bis(2-quinolylmethoxy)phenyl)-4-hydroxy-2-butynoic acid methyl ester,

15 [5,5-bis(4-(2-quinolylmethoxy)phenyl)-5-hydroxy-3-pentyn-1-yl]-2-

iminoxypropionic acid sodium salt,

[5,5-bis(4-(2-quinolylmethoxy)phenyl)-5-hydroxy-3-pentyn-1-yl]-2-

iminoxypropionic acid,

4,4-bis(4-(2-benzoxazolylmethoxy)phenyl)pentanoic acid,

20 4,4-bis(4-(2-pyrimidylmethoxy)phenyl)pentanoic acid,

4,4-bis(4-(4-phenyl-2-thiazolylmethoxy)phenyl)pentanoic acid,

4,4-bis(4-(4-(pyrid-2-yl)-2-thiazolylmethoxy)phenyl)pentanoic acid,

4,4-bis(4-(6-phenyl-2-pyridylmethoxy)phenyl)pentanoic acid,

4,4-bis(4-(5-phenyl-2-pyridylmethoxy)phenyl)pentanoic acid,

25 4,4-bis(4-(6-(pyrid-2-yl)-2-pyridylmethoxy)phenyl)pentanoic acid, and

4,4-bis(4-(4-phenyl-2-pyrimidylmethoxy)phenyl)pentanoic acid.

Preferred compounds of the present invention have the structure defined above wherein Z is selected from the group consisting of (a) COM, (b) CH=N-O-A-COM, (c) CH₂-O-N=A-COM, and (d) OH wherein A is alkylene of one to six carbon atoms, and M is -OH.

More preferred compounds of the present invention have the structure defined immediately above wherein W is selected from the group consisting of (a) quinolyl; (b) quinolyl substituted with a substituent selected from the group consisting of (b-1) halogen, (b-2) alkyl of one to six carbon atoms, (b-3) phenyl, (b-4) phenyl substituted with a substituent selected from the group consisting of (b-4-a) halogen, (b-4-b) alkyl of one to six carbon atoms, (b-4-c) haloalkyl of one to six carbon atoms, and (b-4-d) alkoxy of one to six carbon atoms, (b-5) pyridyl, and (b-6) pyridyl

substituted with a substituent selected from the group consisting of (b-6-a) halogen, (b-6-b) alkyl of one to six carbon atoms, and (b-6-c) alkoxy of one to six carbon atoms; (c) benzothiazolyl; and (d) benzothiazolyl substituted with a substituent selected from the group consisting of (d-1) halogen, (d-2) alkyl of one to six carbon atoms, (d-3) phenyl, (d-4) phenyl substituted with a substituent selected from the group consisting of (d-4-a) halogen, (d-4-b) alkyl of one to six carbon atoms, (d-4-c) haloalkyl of one to six carbon atoms, and (d-4-d) alkoxy of one to six carbon atoms, (d-5) pyridyl, and (d-6) pyridyl substituted with a substituent selected from the group consisting of (d-6-a) halogen, (d-6-b) alkyl of one to six carbon atoms, and (d-6-c) alkoxy of one to six carbon atoms.

Still more preferred compounds have the structure defined immediately above wherein X is alkylene of one to six carbon atoms, and Z is COOH.

The most preferred compounds of the present invention have the structure defined immediately above wherein W is the same at each occurrence and is selected from the group consisting of (a) quinolyl; and (b) quinolyl substituted with a substituent selected from the group consisting of (b-1) halogen, (b-2) alkyl of one to six carbon atoms, (b-3) phenyl, (b-4) phenyl substituted with a substituent selected from the group consisting of (b-4-a) halogen, (b-4-b) alkyl of one to six carbon atoms, (b-4-c) haloalkyl of one to six carbon atoms, and (b-4-d) alkoxy of one to six carbon atoms, (b-5) pyridyl, and (b-6) pyridyl substituted with a substituent selected from the group consisting of (b-6-a) halogen, (b-6-b) alkyl of one to six carbon atoms, and (b-6-c) alkoxy of one to six carbon atoms.

Compounds representative of the most preferred embodiment include, but are not limited to

- 25 4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid,
- 4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid sodium salt,
- 4,4-*bis*(4-(2-quinolylmethoxy)phenyl) pentanoic acid magnesium salt,
- 4,4-*bis*(4-(2-benzothiazolylmethoxy)phenyl)pentanoic acid,
- 4,4-*bis*(4-(2-benzothiazolylmethoxy)phenyl)pentanoic acid sodium salt,
- 30 4,4-*bis*(4-(7-chloro-2-quinolylmethoxy)phenyl)pentanoic acid,
- 4,4-*bis*(4-(7-fluoro-2-quinolylmethoxy)phenyl)pentanoic acid sodium salt,
- 2,2-*bis*(4-(2-quinolylmethoxy)phenyl)propionic acid,
- 3,3-*bis*(4-(2-quinolylmethoxy)phenyl)butanoic acid,
- 5,5-*bis*(4-(2-quinolylmethoxy)phenyl)hexanoic acid,
- 35 5,5-*bis*(4-(2-quinolylmethoxy)phenyl)hexanoic acid sodium salt,
- 4,4-*bis*(3-chloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid,
- 4,4-*bis*(3-chloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid sodium salt,

2,2-bis(4-(2-quinolylmethoxy)phenyl)butyric acid,
2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)propionic acid sodium
salt,
2,2-bis(4-(2-quinolylmethoxy)phenyl)acetic acid,
5 2,2-bis(4-(6-fluoro-2-quinolylmethoxy)phenyl)acetic acid,
2,2-bis(4-(2-quinolylmethoxy)phenyl)propionic acid, and
2,2-bis(4-(2-quinolylmethoxy)phenyl)propionic acid sodium salt.

Lipoxygenase Inhibition Determination

10 Inhibition of leukotriene biosynthesis was evaluated in vitro using an assay involving calcium ionophore-induced LTB₄ expressed in human polymorphonuclear leukocytes (PMNL). Human PMNL isolated from heparinized (20 USP units/mL) venous blood (25 mL) obtained from healthy volunteers was layered over an equal volume of Ficoll-Hypaque Mono-Poly Resolving Medium (ICN Flow, Costa Mesa,
15 CA) and centrifugated at 400 x g for 40 minutes at 20°C. The PMNL was collected, erythrocytes lysed and washed 2x and suspended at 1.0 x 10⁷ cells/mL in Earle's balanced salt solution with 17 mM Earle's HEPES. Aliquots of the cell suspension were preincubated with test compounds dissolved in DMSO (final concentration <2%) for 15 minutes and stimulated with calcium ionophore (final concentration 8.3, µM)
20 for 10 minutes at 37°C. Incubations were stopped with the addition of two volumes of ice-cold methanol followed by centrifuging the cell suspensions at 4°C for 10 minutes at ~450 x g. The amount of LTB₄ in the methanol extract was analyzed by enzyme-linked immunoassay or by HPLC analysis.

The compounds of this invention inhibit leukotriene biosynthesis as shown by
25 the data for representative examples in Table 1.

Table 1

In Vitro Inhibitory Potencies Against 5-Lipoxygenase From
Stimulated LTB₄ Formation in Human Polymorphonuclear Leukocytes

30

Example	IC ₅₀ (µM)
1	0.040
2	0.069
5	0.030
6	0.035
7	0.040

Table 1 (Continued)

Example	IC ₅₀ (μM)
8	0.046
9	0.050
11	0.040
12	0.032
14	0.043
20	0.026
25	0.050
26	0.080
27	0.028
28	0.034
30	0.050
31	0.035
34	0.070
36	0.025
38	0.060

- 5 Inhibition of leukotriene biosynthesis *in vivo* was evaluated using the
 Ionophore A32187-Induced Rat Pleural Inflammation Model. Pleural inflammation
 was induced in male rats following the method of Rao *et al* (Rao, T. S., Currie, J. L.,
 Shaffer, A. F., Isakson, P. C., (1993) Evaluation of 5-lipoxygenase Inhibitors,
 Zileuton, A-78773 and ICI D-2138 in an Ionophore (A-23187) Induced Pleural
 10 Inflammation Model in the Rat, *Life Sciences*, 53: 147 (1993)). Rats were dosed
 with experimental compounds in 0.2% methocel one hour prior to the intrapleural
 injection of the calcium ionophore, A23187. The rats were lightly anesthetized with
 Pentrane (Abbott Laboratories) and injected intrapleurally with 0.5 ml of 2% ethanol
 in injectable saline (Abbott Laboratories) containing 20 μg of A23187 (Cal BioChem-
 15 Novabiochem). Thirty minutes later the animals were euthanized and the pleural
 cavities lavaged with ice cold saline (Abbott Laboratories). The lavage fluid was then
 added to ice cold methanol (final methanol concentration 30%) to lyse cells and
 precipitate protein. Eicosanoids were determined by enzyme immunoassay by
 standard methods.

Table 2
In Vivo Leukotriene Inhibition in Rat Pleural Inflammation

Example	% Inhibition at 3 mg/kg
2	53%
11	42%
14	32%
24	38%
26	50%
27	42%
34	30%
38	40%

5

Pharmaceutical Compositions

The present invention also provides pharmaceutical compositions which comprise compounds of the present invention formulated together with one or more non-toxic pharmaceutically acceptable carriers. The pharmaceutical compositions may be specially formulated for oral administration in solid or liquid form, for parenteral
10 injection, or for rectal administration.

The pharmaceutical compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, or as an oral or nasal spray. The term "parenteral" administration as used herein refers to
15 modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Pharmaceutical compositions of this invention for parenteral injection comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions as well as sterile powders for reconstitution
20 into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by
25 the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

These compositions may also contain adjuvants such as preservative, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents such as sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of the drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or

- a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid,
- b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia,
- c) humectants such as glycerol,
- d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate,

e) solution retarding agents such as paraffin,
f) absorption accelerators such as quaternary ammonium compounds,
g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate,

5 h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

Solid compositions of a similar type may also be employed as fillers in soft
10 and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying
15 agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

The active compounds can also be in micro-encapsulated form, if appropriate,
20 with one or more of the above-mentioned excipients.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and
25 emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

30 Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol
35 and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar, and tragacanth, and mixtures thereof.

Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at room temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

Compounds of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multilamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any nontoxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic.

Methods to form liposomes are known in the art. See, for example, Prescott, Ed., Methods in Cell Biology, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33, *et seq.*

Dosage forms for topical administration of a compound of this invention include powders, sprays, ointments and inhalants. The active compound is mixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives, buffers, or propellants which may be required. Ophthalmic formulations, eye ointments, powders and solutions are also contemplated as being within the scope of this invention.

Actual dosage levels of active ingredients in the pharmaceutical compositions of this invention may be varied so as to obtain an amount of the active compound(s) that is effective to achieve the desired therapeutic response for a particular patient, compositions, and mode of administration. The selected dosage level will depend upon the activity of the particular compound, the route of administration, the severity of the condition being treated, and the condition and prior medical history of the patient being treated. However, it is within the skill of the art to start doses of the compound at levels lower than required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.


Generally dosage levels of about 1 to about 50, more preferably of about 5 to about 20 mg of active compound per kilogram of body weight per day are administered orally to a mammalian patient. If desired, the effective daily dose may be divided into multiple doses for purposes of administration, *e.g.*, two to four separate doses per day.

Preparation Of Compounds Of This Invention

In general, the compounds of this invention are synthesized by reaction schemes 1-3 as illustrated below. The preparation of compounds of this invention wherein Z is CO_2R^6 , wherein R^6 is H or alkyl is outlined in Scheme 1. Reaction of two equivalents of phenol with the requisite carbonyl (keto or aldehyde) ester in the presence of acid gives adducts of formula I (see US Pat. No. 2,933,520). Intermediates of formula I wherein R^6 is H are esterified, for example by reaction with an alcohol in the presence of acid, to give an ester of formula II, which is then reacted with a heteroarylmethyl halide of formula $\text{W-CH}_2\text{X}$ where X is Cl, Br, or I, and W is defined above in the presence of a suitable base such as K_2CO_3 to provide the desired compound III in which R^6 is alkyl. Hydrolysis of the ester, for example using aqueous alkali, provides compounds of formula IV wherein R^6 is H.

15

Scheme 1




A general procedure for the synthesis of compounds of this invention wherein Z is OR^4 or CH=N-O-A-COM is described in Scheme 2. Reduction of ester III, for example using lithium aluminum hydride or sodium borohydride, or acid IV, for example using lithium aluminum hydride or using sodium borohydride to reduce the mixed anhydride made from the acid and ethyl chloroformate, provides alcohol V. Compounds of this invention wherein R^4 is alkyl are then prepared from V using methods well-known in the art such as reaction with alkyl iodide in the presence of base. Alcohol V is converted to compounds of this invention wherein CH=N-O-A-COM by oxidation to the aldehyde, for example using Swern oxidation conditions

(Swern, et al., *J. Org. Chem.*, **1978**, *43*, 2480), followed by reaction with the requisite hydroxylamine derivative $\text{H}_2\text{N-O-A-COM}$.


Scheme 2

5



The preparation of compound of this invention wherein Z is $\text{CH}_2\text{-O-N=A-COM}$ is described in Scheme 3. The hydroxy intermediate V, prepared in Scheme 2 above, is converted to hydroxylamine derivative VIII by known methods such as coupling with N-hydroxyphthalimide under Mitsunobu conditions (triphenylphosphine, diethyl or diisopropylazodicarboxylate; see Mitsunobu, O., *Synthesis*, **1981**, 1), followed by treatment with hydrazine. The hydroxylamine derivative VIII is then reacted with the requisite carbonyl unit, O=A-COM to provide the compounds of this invention represented by the general structure IX.

Scheme 3




The foregoing may be better understood by reference to the following examples which are provided for illustration and are not intended to limit the scope of the invention as it is defined by the appended claims.

5

Example 1

Preparation of 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanoic acid



10

Step 1: 4,4-bis(4-hydroxyphenyl)pentanoic acid methyl ester

To a solution in methanol (120 mL) of 4,4-bis(4-hydroxyphenyl)pentanoic acid (Aldrich Chemical Co., 12 g, 42 mmol) was added concentrated H₂SO₄ (0.5 mL) and the mixture was heated to reflux for 3 hours. After cooling to room temperature the mixture was concentrated *in vacuo* and dissolved in ether (300 mL). The organic layer was washed with saturated aqueous NaHCO₃ (2 x 150 mL) and brine, dried over MgSO₄, filtered, and concentrated *in vacuo* to provide a thick oil which was crystallized from ether/hexane to give 4,4-bis(4-hydroxyphenyl)pentanoic acid methyl ester as an off-white color solid (11.8 g, 94%), mp 130°C.

20

Step 2: 4,4-bis(4-(2-quinolylmethyl)phenyl)pentanoic acid methyl ester

To a solution in dry DMF under N₂ of 4,4-bis(4-hydroxyphenyl)pentanoic acid methyl ester (6.0 g, 20 mmol), prepared as in step 1, was added powdered K₂CO₃ (11.0 g, 80 mmol) and the reaction was stirred for 10 minutes after which 2-chloromethylquinoline hydrochloride (8.5 g, 40 mmol) was added. The mixture was heated at 60 °C for 18 hours and then cooled to room temperature, diluted with EtOAc (200 mL), washed with water and brine, dried over MgSO₄, filtered and concentrated *in vacuo* to provide a residue which was purified by chromatography on silica gel (9:1 CH₂Cl₂/EtOAc) to provide 10.8 (92%) of 4,4-bis(4-(2-quinolylmethyl)phenyl)pentanoic acid methyl ester.

30

Step 3: 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanoic acid

To a solution in 1:1 dioxane/methanol (100 mL) of 4,4-bis(4-(2-quinolylmethyl)phenyl)pentanoic acid methyl ester (3.3 g, 5.7 mmol), prepared as in

step 2, was added aqueous 1N NaOH (10 mL) and the mixture was heated at reflux for 3 hours, cooled to room temperature, concentrated *in vacuo*, diluted with water and neutralized with 10% aqueous citric acid. The solid precipitate was collected by filtration, dried *in vacuo*, and purified by chromatography on silica gel (9:1 CH₂Cl₂/EtOAc, followed by 20:1 CH₂Cl₂/CH₃OH) to provide 2.53 g (70%) of the desired product. Crystallization from methylene chloride-hexanes gave 4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid: mp 105-106°C; ¹H NMR (300 MHz, DMSO-d₆) δ 1.50 (s, 3H), 1.95 (m, 2H), 2.27 (m, 2H), 5.34 (s, 4H), 6.98 (d, 4H, J = 8 Hz), 7.10 (d, 4H, J = 8 Hz), 7.65 (m, 4H), 7.80 (m, 2H), 8.00 (m, 4H), 8.42 (d, 2H, J = 9 Hz), 12.00 (s, 1H); MS (DCI-NH₃) m/e 569 (M+H)⁺. Anal. Calc'd. for C₃₇H₃₂N₂O₄: C, 78.15; H, 5.67; N, 4.93. Found: C, 77.52; H, 5.88; N, 4.60.

Example 2

Preparation of 4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid sodium salt

To a solution in dioxane (10 mL) and methanol (10 mL) of 4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid (320 mg, 0.56 mmol), prepared as in step 1, was added aqueous 1N sodium hydroxide (0.55 mL, 0.55 mmol). The mixture was then concentrated *in vacuo*. The product was crystallized by dissolving in CH₂Cl₂ and precipitation by dropwise addition of a mixture of ethyl acetate-ethyl ether (1:2): ¹H NMR (300 MHz, DMSO-d₆) δ 1.47 (s, 3H), 1.63 (m, 2H), 2.18 (m, 2H), 5.31 (s, 4H), 6.95 (d, 4H, J = 9 Hz), 7.08 (d, 4H, J = 9 Hz), 7.64 (m, 4H), 7.78 (m, 2H), 8.00 (m, 4H), 8.40 (d, 2H, J = 8 Hz); MS (FAB⁺) m/e 591 (M+H)⁺; (FAB⁻) 589 (M-H)⁻. Anal. Calc'd. for C₃₇H₃₁N₂O₄Na•0.25 H₂O: C, 74.67; H, 5.34; N, 4.71; Found: C, 74.57; H, 5.32; N, 4.52.

Example 3


Preparation of 4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid magnesium salt

To a stirred room temperature THF solution of 4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid (0.6 g, 1.06 mmol), prepared as in Example 1, was added MgO (0.021 g, 0.528 mmol). Water was added until the mixture became homogeneous. The reaction was allowed to stir for 24 hours. The solvent was removed *in vacuo* and the residue triturated with hexanes. The precipitate was vacuum filtered and washed with hexanes. The solid was dried *in vacuo* at 55 °C for 48 hours to give 0.510 g (83%) of 4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid magnesium salt as a cream-colored powder: mp 96-110°C; ¹H NMR (300 MHz, DMSO-d₆) δ 1.42 (s, 6H), 1.76 (m, 4H), 2.21 (m, 4H), 5.27 (s, 8H), 6.91 (d, 8H, J = 7.5 Hz), 7.03 (d, 8H, J = 7.5 Hz), 7.58 (m, 8H), 7.74 (t, 4H, J = 7.5 Hz), 7.96

(m, 8H), 8.34 (m, 4H). Anal. Calc'd. for $C_{74}H_{62}N_4O_8Mg \cdot 1.50 H_2O$: C, 74.90; H, 5.52; N, 4.72. Found: C, 74.81; H, 5.64; N, 4.68.

Example 4


5 Preparation of 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentan-1-ol



To a mixture in THF (50 mL) of 4,4-bis(4-(2-quinolylmethoxy)phenyl)-
 10 pentanoic acid 2-quinolylmethyl ester (2.1 g, 3 mmol), prepared as in Example 1, and sodium borohydride (380 mg, 10 mmol) was added dropwise methanol at 50-55 °C and the mixture was stirred for 30 minutes. The mixture was cooled to room temperature, poured into water (50 mL) and acidified to pH 4. The resulting mixture was extracted with ethyl acetate, dried over $MgSO_4$, filtered, and concentrated *in*
 15 *vacuo*. The residue was purified by chromatography on silica gel (methylene chloride-ethyl acetate 3:1) to afford 1.46 g (88%) of 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentan-1-ol: 1H NMR (300 MHz, $DMSO-d_6$) δ 1.18 (m, 2H), 1.48 (s, 3H), 2.00 (m, 2H), 3.34 (m, 2H), 4.34 (t, 1H, $J = 6$ Hz), 5.30 (s, 4H), 6.96 (d, 4H, $J = 9$ Hz), 7.09 (d, 4H, $J = 9$ Hz), 7.64 (m, 4H), 7.79 (m, 2H), 8.00 (t, 4H, $J = 8$ Hz), 8.41 (d, 2H, $J = 8$ Hz); MS (DCI- NH_3) m/e 555 ($M+H$) $^+$. Anal. Calc'd. for $C_{37}H_{34}N_2O_3 \cdot 0.5 H_2O$: C, 78.84; H, 6.44; N, 4.97. Found: C, 78.67; H, 5.95; N, 4.70.

Example 5

25 Preparation of [4,4-bis(4-(2-quinolylmethoxy)phenyl)pent-1-yl]liminooxyacetic acid



Step 1: *N*-phthaloyl-*O*-[4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pent-1-yl]hydroxylamine


To a solution in THF (35 mL) of 4,4-*bis*(4-(2-quinolylmethoxy)phenyl)-
 pentan-1-ol (1.11 g, 2 mmol), prepared as in Example 4, *N*-hydroxyphthalimide (326
 5 mg, 2 mmol) and triphenylphosphine (786 mg, 3 mmol) was added a solution of
 DIAD (0.6 mL, 3 mmol) in THF (5 mL), and the resulting mixture was stirred at room
 temperature for 10 hours. The reaction mixture was concentrated *in vacuo* and
 purified by chromatography on silica gel (methylene chloride-ethyl acetate 12:1) to
 provide 1.33 g of *N*-phthaloyl-*O*-[4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pent-1-
 10 yl]hydroxylamine.

Step 2: *O*-[4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pent-1-yl]hydroxylamine

To a solution in dioxane (15 mL) and ethanol (15 mL) of the *N*-phthaloyl-*O*-
 [4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pent-1-yl]hydroxylamine prepared in step 1
 15 was added hydrazine hydrate (0.25 mL, 4 mmol) and the mixture was heated at reflux
 for 30 minutes. The mixture was then treated with aqueous 10% sodium carbonate
 (20 mL) and extracted with ethyl acetate. The organic extract was washed with water
 and brine, dried over MgSO₄, filtered, and concentrated *in vacuo*. The residue was
 purified by chromatography on silica gel (ethyl acetate) to afford 420 mg of *O*-[4,4-
 20 *bis*(4-(2-quinolylmethoxy)phenyl)pent-1-yl]hydroxylamine.

Step 3: [4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pent-1-yl]iminoxyacetic acid

A solution in dioxane (10 mL), methanol (5 mL) and water (2 mL) of *O*-[4,4-
bis(4-(2-quinolylmethoxy)phenyl)pent-1-yl]hydroxylamine (114 mg, 0.2 mmol),
 25 prepared as in step 2, glyoxylic acid (19 mg, 0.2 mmol) and acetic acid (0.012 mL,
 0.2 mmol) was stirred at ambient temperature for 16 hours. The mixture was then
 concentrated *in vacuo* and the residue was dissolved in ethyl acetate (50 mL). The
 resulting solution was washed with water and brine, dried over MgSO₄, filtered, and
 concentrated *in vacuo*. The residue was crystallized from methylene chloride-hexane
 30 to afford 108 mg (86%) of [4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pent-1-
 yl]iminoxyacetic acid: mp 80-82°C; ¹H NMR (300 MHz, DMSO-d₆) δ 1.40 (m,
 2H), 1.53 (s, 3H), 2.04 (m, 2H), 4.10 (t, 2H, J = 7 Hz), 5.31 (s, 4H), 6.95 (d, 4H,
 J = 9 Hz), 7.08 (d, 4H, J = 9 Hz), 7.52 (s, 1H), 7.63 (m, 4H), 7.78 (m, 2H), 8.00
 (m, 4H), 8.40 (d, 2H, J = 8 Hz); MS (DCI-NH₃) m/e: 626 (M+H)⁺. Anal. Calc'd.
 35 for C₃₉H₃₅N₃O₅: C, 74.86; H, 5.64; N, 6.72; Found: C, 74.89; H, 6.03; N, 6.42.

Example 6**Preparation of [4,4-bis(4-(2-quinolylmethoxy)phenyl)pent-1-yl]-2-iminoxypropionic acid**

5

Step 1: [4,4-bis(4-(2-quinolylmethoxy)phenyl)pent-1-yl]-2-iminoxypropionic acid methyl ester


A mixture in dioxane (20 mL) and methanol (20 mL) of *O*-[4,4-bis(4-(2-quinolylmethoxy)phenyl)pent-1-yl]hydroxylamine (306 mg, 0.54 mmol), prepared as in Example 5, step 2, methyl pyruvate (0.055 mL, 0.55 mmol) and acetic acid (0.033 mL, 0.55 mmol) was stirred at room temperature for 12 hours. The mixture was then partitioned between aqueous 10% sodium bicarbonate and ethyl acetate. The ethyl acetate extract was washed with water and brine, dried over MgSO₄, filtered, and concentrated *in vacuo*. The residue was purified by chromatography on silica gel (methylene chloride-ethyl acetate 4:1) to afford 320 mg of [4,4-bis(4-(2-quinolyl-methoxy)phenyl)pent-1-yl]-2-iminoxypropionic acid methyl ester.

Step 2: [4,4-bis(4-(2-quinolylmethoxy)phenyl)pent-1-yl]-2-iminoxypropionic acid

The desired compound was prepared according to the method of Example 1, step 3, except substituting [4,4-bis(4-(2-quinolylmethoxy)phenyl)pent-1-yl]-2-iminoxypropionic acid methyl ester, prepared as in step 1, for 4,4-bis(4-(2-quinolylmethyl)phenyl)pentanoic acid methyl ester: mp 80-82°C; ¹H NMR (300 MHz, DMSO-d₆) δ 1.42 (m, 2H), 1.53 (s, 3H), 1.90 (s, 3H), 2.05 (m, 2H), 4.11 (t, 2H, J = 7 Hz), 5.33 (s, 4H), 6.95 (d, 4H, J = 9 Hz), 7.09 (d, 4H, J = 9 Hz), 7.64 (m, 4H), 7.78 (m, 2H), 8.00 (m, 4H), 8.41 (d, 2H, J = 8 Hz); MS (DCI-NH₃) m/e 640 (M+H)⁺. Anal. Calc'd. for C₄₀H₃₇N₃O₅: C, 75.10; H, 5.83; N, 6.57. Found: C, 74.86; H, 6.11; N, 6.27.

Example 7

Preparation of [4,4-bis(4-(2-quinolylmethoxy)phenyl)pent-1-yl]oximinoacetic acid



5


Step 1: 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanal

To a solution in DMSO (20 mL) of 4,4-bis(4-(2-quinolylmethoxy)phenyl)-
 pentan-1-ol (410 mg, 0.74 mmol), prepared as in Example 4, and 1,3-dicyclo-
 hexylcarbodiimide (515 mg, 2.5 mmol) was added aqueous 1M phosphoric acid
 10 (0.5 mL) and the resulting mixture was stirred at room temperature for 4 hours. Ethyl
 acetate (80 mL) was added and dicyclohexylurea was filtered off. The filtrate was
 washed with water and brine dried over MgSO₄, filtered, and concentrated *in vacuo*.
 The residue was purified by chromatography on silica gel (methylene chloride-ethyl
 acetate 9:1) to afford 280 mg of 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanal.

15

Step 2: [4,4-bis(4-(2-quinolylmethoxy)phenyl)pent-1-yl]oximinoacetic acid

A mixture in dioxane (10 mL), methanol (10 mL) and water (5 mL) of the
 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanal (280 mg, 0.5 mmol) prepared in step
 1, carboxymethoxylamine hemihydrochloride (110 mg, 0.5 mmol), and sodium
 20 acetate trihydrate (69 mg, 0.5 mmol) was stirred at ambient temperature for 12 hours.
 The mixture was diluted with water and extracted with ethyl acetate. The ethyl acetate
 extract was washed with water and brine, dried over MgSO₄, filtered, and
 concentrated *in vacuo*. The residue was recrystallized from methylene chloride-
 hexane to afford 250 mg (80 %) of [4,4-bis(4-(2-quinolylmethoxy)phenyl)pent-1-
 25 yl]oximinoacetic acid: mp 78-80°C; ¹H NMR (300 MHz, DMSO-d₆) δ 1.53 (s, 3H),
 1.88 (m 1H), 2.05 (m, 1H), 2.22 (m, 2H), 4.40 and 4.45 (two s, 1:1, 2H), 5.34 (s,
 4H), 6.75 and 7.45 (two t, 1:1, 1H), 6.96 (d, 4H, J = 9 Hz), 7.10 (dd, 4H, J = 9, 7
 Hz), 7.64 (m, 4H), 7.78 (m, 2H), 8.00 (m, 4H), 8.41 (d, 2H, J = 8 Hz), 12.62 (bs
 , 1H); MS (DCI-NH₃) m/e 626 (M+H)⁺. Anal. Calc'd. for C₃₉H₃₅N₃O₅•0.5 H₂O:
 30 C, 73.79; H, 5.72; N, 6.62. Found: C, 73.94; H, 6.13; N, 6.68.

Example 8**Preparation of 4,4-bis(4-(2-benzothiazolymethoxy)phenyl)pentanoic acid**

5

The desired product was prepared according to the procedure of Example 1, except substituting 2-chloromethylbenzothiazole for 2-chloromethylquinoline: mp 185-186°C; ¹H NMR (300 MHz, DMSO-d₆) δ 1.60 (s, 3H), 1.94 (m, 2H), 2.29 (m, 2H), 5.05 (s, 4H), 7.01 (d, 4H, J = 9 Hz), 7.13 (d, 4H, J = 9 Hz), 7.50 (m, 4H), 8.03 (d, 2H, J = 9 Hz), 8.13 (m, 2H), 12.04 (bs, 1 H); MS (DCI-NH₃) m/e 598 (M+NH₄)⁺, 581 (M+H)⁺. Anal. Calc'd. for C₃₃H₂₈N₂O₄S₂: C, 68.27; H, 4.86; N, 4.83. Found C, 68.06; H, 4.70; N, 4.64.


10

Example 9**Preparation of 4,4-bis(4-(2-benzothiazolymethoxy)phenyl)pentanoic acid sodium salt**

The desired salt was prepared according to the procedure of Example 2, except substituting 4,4-bis(4-(2-benzothiazolymethoxy)phenyl)pentanoic acid, prepared as in Example 8, for 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanoic acid: mp 105-108°C; ¹H NMR (300 MHz, DMSO-d₆) δ 1.48 (s, 3H), 1.58 (m, 2H), 2.19 (m, 2H), 5.54 (s, 4H), 6.98 (d, 4H, J = 9 Hz), 7.10 (d, 4H, J = 9 Hz), 7.5 (m, 4H), 8.01 (d, 2H, J = 9 Hz), 8.11 (d, 2H, J = 9 Hz). MS (FAB) m/e 625 (M+Na)⁺, 603 (M+H)⁺. Anal. Calc'd. for C₃₃H₂₇N₂O₄SN_a•1.5 H₂O: C, 62.95; H, 4.80; N, 4.45. Found: C, 63.11; H, 4.72; N, 4.26.

20

25

Example 10**Preparation of 4,4-bis(4-(7-chloro-2-quinolylmethoxy)phenyl)pentanoic acid**


30

The desired compound was prepared according to the procedure of Example 1, except substituting 2-chloromethyl-7-chloroquinoline for 2-chloromethylquinoline:

- mp 88-90°C; ^1H NMR (300 MHz, DMSO- d_6) δ 1.51 (s, 3H), 1.95 (m, 2H), 2.28 (m, 2H), 5.33 (s, 4H), 6.97 (d, 4H, $J = 9$ Hz), 7.09 (d, 4H, $J = 9$ Hz), 7.66 (dd, 2H, $J = 9, 2$ Hz), 7.71 (d, 2H, $J = 9$ Hz), 8.06 (m, 4H), 8.47 (d, 2H, $J = 9$ Hz), 12.03 (bs, 1H); MS (DCI- NH_3) m/e 637 ($\text{M}+\text{H}$) $^+$. Anal. Calc'd. for $\text{C}_{37}\text{H}_{30}\text{N}_2\text{O}_4$: C, 69.81; H, 4.75; N, 4.40. Found: C, 69.77; H, 5.05; N, 4.17.

Example 11

Preparation of 4,4-bis(4-(6-fluoro-2-quinolylmethoxy)phenyl)pentanoic acid sodium salt



Step 1: 4,4-bis(4-(6-fluoro-2-quinolylmethoxy)phenyl)pentanoic acid


- The desired compound was prepared according to the procedure of Example 1, except substituting 2-chloromethyl-6-fluoroquinoline for 2-chloromethylquinoline.

Step 2: 4,4-bis(4-(6-fluoro-2-quinolylmethoxy)phenyl)pentanoic acid sodium salt

- The desired salt was prepared according to the procedure of Example 2, except substituting 4,4-bis(4-(6-fluoro-2-quinolylmethoxy)phenyl)pentanoic acid, prepared as in step 1, for 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanoic acid: mp 97-99°C; ^1H NMR (300 MHz, DMSO- d_6) δ 1.48 (s, 3H), 1.70 (m, 2H), 2.1 (m, 2H), 5.31 (s, 4H), 6.95 (d, 4H, $J = 9$ Hz), 7.08 (d, 4H, $J = 9$ Hz), 7.68 (m, 4H), 7.80 (dd, 2H, $J = 9, 3$ Hz), 8.08 (dd, 2H, $J = 9, 6$ Hz), 8.41 (d, 2H); MS (FAB) m/e 627 ($\text{M}+\text{Na}$) $^+$, 605 ($\text{M}+\text{H}$) $^+$. Anal. Calc'd. for: $\text{C}_{37}\text{H}_{29}\text{F}_2\text{N}_2\text{O}_4\text{Na}$: C, 70.92; H, 4.76; N, 4.47. Found: C, 70.79; H, 4.84; N, 4.30.

Example 12

Preparation of 2,2-bis(4-(2-quinolylmethoxy)phenyl)propionic acid



Step 1: 2,2-bis(4-hydroxyphenyl)propionic acid


To a cooled mixture of phenol (9.4 g, 0.1 mol), pyruvic acid (4.4 g, 0.05 mol), and water was added concentrated H₂SO₄ (4.5 mL, 18.0 g) dropwise with stirring. After 15 minutes the ice bath was removed and the reaction was warmed to room temperature and stirred for 18 hours. The reaction mixture was diluted with ethyl ether/water (200 ml 1:1) and the layers were separated. The organic layer was extracted with saturated aqueous NaHCO₃. The aqueous extract was then acidified, extracted with ethyl ether, dried over MgSO₄, filtered, and concentrated *in vacuo* to give 2,2-bis(4-hydroxyphenyl)propionic acid (6.2 g) as a colorless sticky oil.

Step 2: 2,2-bis(4-(2-quinolylmethoxy)phenyl)propionic acid


The desired compound was prepared according to the method of Example 1, except substituting 2,2-bis(4-hydroxyphenyl)propionic acid, prepared as in step 1, for 4,4-bis(4-hydroxyphenyl)pentanoic acid: mp 208-210°C; ¹H NMR (300 MHz, DMSO-d₆) δ 1.75 (s, 3H), 5.33 (s, 4H), 7.01 (d, 4H, J = 9 Hz), 7.12 (d, 4H, J = 9 Hz), 7.63 (m, 4H), 7.78 (m, 2H), 8.01 (t, 4H, J = 8 Hz), 8.42 (d, 2H, J = 8 Hz), 12.65 (bs, 1 H); MS (DCI-NH₃) m/e 541 (M+H)⁺. Anal. Calc'd. for C₃₅H₂₈N₂O₄•H₂O: C, 75.19; H, 5.28; N, 5.02. Found: C, 74.85; H, 4.89; N, 4.92.

Example 13

Preparation of 3,3-bis(4-(2-quinolylmethoxy)phenyl)butanoic acid



The desired product was prepared according to the procedure of Example 12, except substituting ethyl acetoacetate for pyruvic acid and hydrolysis of the intermediate ethyl ester as described in Example 1, step 3: mp 94-96°C; ¹H NMR (300 MHz, DMSO-d₆) δ 1.75 (s, 3H), 3.02 (s, 2H), 5.32 (s, 4H), 6.95 (d, 4H, J = 9 Hz), 7.11 (d, 4H, J = 9 Hz), 7.65 (m, 4H), 7.78 (m, 2H), 8.01 (t, 4H, J = 8 Hz), 8.41 (d, 2H, J = 8 Hz), 11.83 (bs, 1H); MS (DCI-NH₃) m/e 555 (M+H)⁺. Anal. Calc'd. for C₃₆H₃₀N₂O₄•0.5H₂O: C, 76.71; H, 5.54; N, 4.95. Found: C, 76.23; H, 5.43; N, 4.66.

Example 14**Preparation of 5,5-bis(4-(2-quinolylmethoxy)phenyl)hexanoic acid**

5

The desired compound was prepared according to the procedure of Example 12, except substituting 4-acetyl butyric acid for pyruvic acid: mp 87-89°C; ¹H NMR (300 MHz, DMSO-d₆) δ 1.25 (m, 2H), 1.51 (s, 3H), 2.0 (m, 2H), 2.17 (t, 2H, J = 8 Hz), 5.32 (s, 4H), 6.96 (d, 4H, J = 9 Hz), 7.09 (d, 4H, J = 9 Hz), 7.62 (m, 4H), 7.68 (d, 2H, J = 9 Hz), 7.78 (m, 2H), 8.02 (t, 2H, J = 8 Hz), 8.41 (d, 2H, J = 8 Hz), 11.97 (bs, 1H); MS (FAB) m/e 583 (M+H)⁺. Anal. Calc'd. for C₃₈H₃₄N₂O₄: C, 78.34; H, 5.87; N, 4.81. Found: C, 77.97; H, 6.0; N, 4.63.


10

Example 15**Preparation of 5,5-bis(4-(2-quinolylmethoxy)phenyl)hexanoic acid sodium salt**

The desired compound was prepared according to the method of Example 2, except substituting 5,5-bis(4-(2-quinolylmethoxy)phenyl)hexanoic acid, prepared as in Example 14, for 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanoic acid: ¹H NMR (300 MHz, DMSO-d₆) δ 1.20 (m, 2H), 1.51 (s, 3H), 1.81 (t, 2H, J = 8 Hz), 1.95 (m, 2H), 5.32 (s, 4H), 6.94 (d, 4H, J = 9 Hz), 7.09 (d, 4H, J = 9 Hz), 7.61 (t, 2H, J = 8 Hz), 7.68 (d, 2H, J = 8 Hz), 7.80 (m, 2H), 8.01 (t, 4H, J = 8 Hz), 8.42 (d, 2H, J = 8 Hz). MS (FAB) m/e 605 (M+Na)⁺, 583 (M+H)⁺. Anal. Calc'd. for C₃₈H₃₃N₂O₄Na • 0.5 H₂O: C, 74.39; H, 5.55; N, 4.57. Found: C, 74.64; H, 5.64; N, 4.36.

20

25


Example 16**Preparation of 4,4-bis-(4-(2-pyridylmethoxy)phenyl)pentanoic acid sodium salt**

Step 1: 4,4-bis-(4-(2-pyridylmethoxy)phenyl)pentanoic acid

The desired compound was prepared according to the procedure of Example 1, except substituting 2-picolyl chloride for 2-chloromethylquinoline.

5 Step 2: 4,4-bis-(4-(2-pyridylmethoxy)phenyl)pentanoic acid sodium salt


The desired compound was prepared according to Example 2, except substituting 4,4-bis-(4-(2-pyridylmethoxy)phenyl)pentanoic acid, prepared as in step 1, for 4,4-bis-(4-(2-quinolylmethoxy)phenyl)pentanoic acid: ¹H NMR (300 MHz, DMSO-d₆) δ 1.49 (s, 3H), 1.53 (m, 2H), 2.21 (m, 2H), 5.13 (s, 4H), 6.92 (d, 4H, J = 9 Hz), 7.08 (d, 4H, J = 9 Hz), 7.34 (m, 2H), 7.51 (d, 2H, J = 9 Hz), 7.34 (m, 2H), 7.51 (d, 2H, J = 9 Hz), 7.84 (dt, 2H, J = 9, 2 Hz), 8.56 (d, 2H, J = 4.5 Hz); MS (FAB) m/e 491 (M+Na)⁺, 469 (M+H)⁺. Anal. Calc'd. for C₂₉H₂₇N₂O₄Na•0.5H₂O: C, 69.72; H, 5.57; N, 5.57. Found: C, 69.45; H, 5.59; N, 5.29.

15 **Example 17**Preparation of 4,4-bis-(4-(2-quinoxylmethoxy)phenyl)pentanoic acid sodium salt

20

The desired compound was prepared according to Example 16, except substituting 2-chloromethylquinoxaline for 2-picolyl chloride: ¹H NMR (300 MHz, DMSO-d₆) δ 1.49 (s, 3H), 1.66 (m, 2H), 2.22 (m, 2H), 5.42 (s, 4H), 6.99 (d, 4H, J = 9 Hz), 7.11 (d, 4H, J = 9 Hz), 7.88 (m, 4H), 8.11 (m, 4H), 9.11 (s, 2H); MS (FAB) m/e 593 (M+Na)⁺, 571 (M+H)⁺. Anal. Calc'd. for C₃₅H₂₉N₄O₄Na•H₂O: C, 68.91; H, 5.11; N, 9.19. Found: C, 68.58, H, 5.15, N, 8.99.

25

Example 18**Preparation of 4,4-bis(4-(1-methyl-2-benzimidazolylmethoxy)phenyl)pentanoic acid**

5


The desired product was prepared according to the procedure of Example 1, except substituting 1-methyl-2-chloromethylbenzimidazole for 2-chloromethyl-quinoline: mp 110-112°C; ^1H NMR (300 MHz; DMSO- d_6) δ 1.52 (s, 3H), 1.95 (m, 2H), 2.28 (m, 2H), 3.85 (s, 6H), 5.37 (s, 4H), 7.02 (d, 4H, $J = 9$ Hz), 7.10 (d, 4H, $J = 9$ Hz), 7.35 (m, 4H), 7.56 (m, 2H), 7.64 (m, 2H), 12.00 (bs, 1 H); MS (FAB+) m/e 575 ($M+H$) $^+$; (FAB-) m/e 573 ($M-H$) $^-$. Anal. Calc'd. for $C_{35}H_{34}N_4O_4 \cdot H_2O$: C, 70.93; H, 6.12; N, 9.45. Found: C, 70.79; H, 6.11; N, 8.87.

15

Example 19**Preparation of 4,4-bis(4-(1-methyl-2-benzimidazolylmethoxy)phenyl)pentanoic acid sodium salt**

The desired product was prepared according to the method of Example 2, except substituting 4,4-bis(4-(1-methyl-2-benzimidazolylmethoxy)phenyl)pentanoic acid, prepared as in Example 18, for 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanoic acid: ^1H NMR (300 MHz, DMSO- d_6) δ 1.48 (s, 3H), 1.60 (m, 2H), 2.20 (m, 2H), 3.85 (s, 6H), 5.35 (s, 4H), 7.00 (d, 4H, $J = 9$ Hz), 7.09 (d, 4H, $J = 9$ Hz), 7.35 (m, 4H), 7.56 (m, 2H), 7.64 (m, 2H); MS (FAB+) m/e 597 ($M+Na$) $^+$, 575 ($M+H$) $^+$, (FAB-) m/e 573 ($M-H$) $^-$. Anal. Calc'd. for $C_{35}H_{33}N_4O_4Na \cdot 1.5H_2O$: C, 67.53; H, 5.81; N, 9.00. Found: C, 67.50; H, 5.85; N, 8.56.

30

Example 20**Preparation of 4,4-bis(3-chloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid**

Step 1: 4,4-bis(3-chloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid methyl ester

A mixture of 4,4-bis(4-hydroxyphenyl)pentanoic acid (5.72 g, 20 mmol), and N-chlorosuccinimide (5.84 g, 44 mmol) in chloroform (120 mL) and dioxane (30 mL) was heated at reflux for 5 hours. The reaction mixture was concentrated *in vacuo* and the residue was dissolved in methanol (100 mL). The methanol solution was cooled to -70 °C, thionyl chloride (3 mL) was added, and the mixture was left at ambient temperature for 16 hours. The methanol was removed *in vacuo* and DMF (150 mL), potassium carbonate (13.8 g, 100 mmol) and 2-chloromethyl-quinoline hydrochloride (9 g, 42 mmol) were added to the residue. The resulting mixture was stirred at room temperature for 10 hours. The mixture was diluted with water (400 mL) and extracted with ethyl acetate. The organic layer was washed with water and brine, dried over MgSO₄, filtered, and concentrated *in vacuo*. The residue was purified by chromatography on silica gel (methylene chloride-ethyl acetate 15:1) to afford 8 g of 4,4-bis(3-chloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid methyl ester and 1.2 g of 4-(3-chloro-4-(2-quinolylmethoxy)phenyl)-4-(3,5-dichloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid methyl ester.

Step 2: 4,4-bis(3-chloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid

The desired compound was prepared according to the method of Example 1, step 3, except substituting 4,4-bis(3-chloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid methyl ester, prepared as in step 1, for 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanoic acid methyl ester: mp 91-94°C; ¹H NMR (300 MHz, DMSO-d₆) δ 1.53 (s, 3H), 1.96 (m, 2H), 2.30 (m, 2H), 5.44 (s, 4H), 7.09 (dd, 2H, J = 3, 9 Hz), 7.20 (d, 2H, J = 9 Hz), 7.24 (d, 2H, J = 3 Hz), 7.62 (m, 2H), 7.71 (d, 2H, J = 9 Hz), 7.80 (m, 2H), 8.00 (m, 4H), 8.44 (d, 2H, J = 9 Hz), 12.08 (bs, 1H); MS (DCI-NH₃) m/e 637 (M+H)⁺. Anal. Calc'd. for C₃₇H₃₀Cl₂N₂O₄ x H₂O: C, 67.79; H, 4.92; N, 4.27. Found: C, 68.02; H, 4.85; N, 3.94.

Example 21

Preparation of 4,4-bis(3-chloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid sodium salt


The desired product was prepared according to the method of Example 2, except substituting 4,4-bis(3-chloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid, prepared as in Example 20, for 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanoic acid: ¹H NMR (300 MHz, DMSO-d₆) δ 1.47 (s, 3H), 1.85 (m, 2H), 2.25 (m, 2H), 5.43 (s, 4H), 7.06 (dd, 2H, J = 3, J = 9 Hz), 7.18 (d, 2H, J = 9 Hz), 7.22 (d, 2H, J = 3 Hz), 7.61 (m, 2H), 7.71 (d, 2H, J = 9 Hz), 7.79 (m, 2H), 8.00 (m, 4H), 8.42 (d,

2H, J = 9 Hz); MS (FAB+) m/e 659 (M+Na)⁺, 637 (M+H)⁺, MS (FAB-) m/e 635 (M-H)⁻. Anal. Calc'd. for C₃₇H₂₉Cl₂N₂O₄Na: C, 67.38; H, 4.43; N, 4.25. Found: C, 67.74; H, 4.89; N, 3.96.

5

Example 22

Preparation of [4,4-bis(3-chloro-4-(2-quinolylmethoxy)phenyl)pent-1-yl]-2-iminoxypropionic acid sodium salt



10

Step 1: 4,4-bis(3-chloro-4-(2-quinolylmethoxy)phenyl)pentan-1-ol

The desired compound was prepared according to the method of Example 4, except substituting 4,4-Bis(3-chloro-4-hydroxyphenyl)pentanoic acid, prepared as in Example 20, for 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanoic acid.

15

Step 2: O-[4,4-bis(3-chloro-4-(2-quinolylmethoxy)phenyl)pent-1-yl]hydroxylamine

The desired compound was prepared according to the method of Example 5, steps 1 and 2, except substituting 4,4-bis(3-chloro-4-(2-quinolylmethoxy)phenyl)pentan-1-ol, prepared as in step 1, for 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentan-1-ol.

20

Step 3: [4,4-bis(3-chloro-4-(2-quinolylmethoxy)phenyl)pent-1-yl]-2-iminoxypropionic acid

The desired compound was prepared according to the method of Example 6, except substituting O-[4,4-bis(3-chloro-4-(2-quinolylmethoxy)phenyl)pent-1-yl]hydroxylamine, prepared as in step 2, for O-[4,4-bis(4-(2-quinolylmethoxy)phenyl)pent-1-yl]hydroxylamine.

25

Step 4: [4,4-bis(3-chloro-4-(2-quinolylmethoxy)phenyl)pent-1-yl]-2-iminoxypropionic acid sodium salt


30

The desired compound was prepared according to the method of Example 2, except substituting [4,4-bis(3-chloro-4-(2-quinolylmethoxy)phenyl)pent-1-yl]-2-iminoxypropionic acid, prepared as in step 3, for 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanoic acid: ¹H NMR (300 MHz, DMSO-d₆) δ 1.31 (m,

2H), 1.54 (s, 3H), 1.79 (s, 3H), 2.04 (m, 2H), 3.93 (t, 2H, $J = 7$ Hz), 5.42 (s, 4H), 7.09 (dd, 2H, $J = 3, 9$ Hz), 7.20 (m, 4H), 7.62 (m, 2H), 7.71 (d, 2H, $J = 8$ Hz), 7.80 (m, 2H), 8.00 (m, 4H), 8.44 (d, 2H, $J = 8$ Hz); MS (FAB+) m/e 730 ($M+Na$)⁺, 708 ($M+H$)⁺, (FAB-) m/e 706 ($M-H$)⁻. Anal. Calc'd. for C₄₀H₃₄N₃Cl₂O₅Na•H₂O: C, 64.17; H, 4.84; N, 5.61. Found: C, 64.40; H, 4.87; N, 5.37.

Example 23


Preparation of 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanoic acid *N,N*-diethylhydroxylamine ester



To a solution in methylene chloride (10 mL) of 4,4-bis(4-(2-quinolylmethoxy)phenyl) pentanoic acid (570 mg, 1 mmol), prepared as in Example 1, was added 1,1'-carbonyldiimidazole (162 mg, 1.1 mmol) and the resulting mixture was stirred at room temperature for 25 minutes. *N,N*-Diethylhydroxylamine (0.14 mL, 1.2 mmol) was then added and stirring was continued for additional 30 minutes. The solution was concentrated *in vacuo* and the residue was purified by chromatography on silica gel (methylene chloride-ethyl acetate 4:1) to afford 420 mg (66 %) of 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanoic acid *N,N*-diethylhydroxylamine ester: ¹H NMR (300 MHz, DMSO-d₆) δ 0.92 (t, 6H, $J = 7$ Hz), 1.51 (s, 3H), 2.03 (m, 2H), 2.30 (m, 2H), 2.75 (q, 4H, $J = 7$ Hz), 5.32 (s, 4H), 6.98 (d, 4H, $J = 9$ Hz), 7.09 (d, 4H, $J = 9$ Hz), 7.65 (m, 4H), 7.78 (m, 2H), 8.00 (m, 4H), 8.41 (d, 2H, $J = 9$ Hz); MS (DCI-NH₃) m/e 640 ($M+H$)⁺. Anal. Calc'd. for C₄₁H₄₁N₃O₄: C, 76.97; H, 6.46; N, 6.57. Found: C, 76.82; H, 6.49; N, 6.50.


Example 24

Preparation of 2,2-bis(4-(2-quinolylmethoxy)phenyl)butyric acid



The desired product was prepared according to the procedure of Example 12, except substituting 2-ketobutyric acid for pyruvic acid: mp 83-92°C; ¹H NMR (300 MHz, DMSO-d₆) δ 0.65 (m, 3H), 2.25 (m, 2H), 5.35 (s, 4H), 7.00 (d, 4H, J = 9 Hz), 7.16 (d, 4H, J = 9 Hz), 7.63 (m, 2H), 7.68 (d, 2H, J = 9 Hz), 7.79 (m, 2H), 8.01 (m, 4H), 8.42 (d, 2H, J = 9 Hz), 12.54 (br s, 1H); MS (DCI-NH₃) m/e 555 (M+H)⁺. Anal. Calc'd. for C₃₆H₃₀N₂O₄•0.55 H₂O: C, 76.59; H, 5.55; N, 4.96. Found: C, 76.90; H, 5.78; N, 4.57.

10

Example 25Preparation of 1,1-bis(4-(2-quinolylmethoxy)phenyl)ethanol15 Step 1: bis(4-(2-quinolylmethoxy)phenyl) ketone

To a solution of 4,4'-dihydroxybenzophenone (4.22 g, 20 mmol) and K₂CO₃ (16.5 g, 120 mmol) in DMF (75 mL) was added 2-chloromethylquinoline hydrochloride (8.56 g, 40 mmol) and the resulting solution was stirred at 60 °C or 16 hours. The reaction mixture was then poured into ice water (100 mL) and the resulting solid was collected by filtration, slurried in 20% ether/hexane, filtered, and dried *in vacuo* to afford bis(4-(2-quinolylmethoxy)phenyl) ketone (9.3 g, 94%) as white solid.

25 Step 2: 1,1-bis(4-(2-quinolylmethoxy)phenyl)ethanol


To a -78 °C solution in THF (20 mL) of bis(4-(2-quinolylmethoxy)phenyl) ketone (992 mg, 2 mmol), prepared as in step 1, was added methylmagnesium bromide (3M solution in ethyl ether, 0.8 mL, 2.4 mmol) and the resulting mixture was stirred at room temperature for 12 hours. The mixture was then quenched with saturated aqueous ammonium chloride and extracted with ethyl acetate. The organic phase was washed with water and brine, dried over MgSO₄, filtered, and concentrated *in vacuo*. The residue was purified by chromatography on silica gel (methylene chloride-ethyl acetate 4:1) to afford 920 mg (90%) of 1,1-bis(4-(2-quinolylmethoxy)phenyl)ethanol: mp 129-131°C; ¹H NMR (300 MHz, DMSO-d₆) δ 1.74 (s, 3H), 5.32 (s, 4H), 5.48 (s, 1H), 6.95 (d, 4H, J = 9 Hz), 7.30 (d, 4H, J = 9

Hz), 7.63 (m, 4H), 7.78 (m, 2H), 8.01 (m, 4H), 8.39 (d, 2H, $J = 8$ Hz); MS (DCI- NH_3) m/e 513 ($\text{M}+\text{H}$) $^+$. Anal. Calc'd. for $\text{C}_{34}\text{H}_{28}\text{N}_2\text{O}_3$: C, 79.67; H, 5.51; N, 5.46. Found: C, 79.48; H, 5.62; N, 5.25.

5

Example 26

Preparation of 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)propionic acid sodium salt



10

Step 1: 2,2-dimethyl-3,3-bis(4-hydroxyphenyl)propionic acid methyl ester

To a stirred 0 °C solution in methanol (2 mL) of phenol (3.76 g, 40 mmol) and methyl-3-carboxaldehyde-2,2-dimethyl propionate (2.6 g, 20 mmol) was added dropwise 10 g of sulfuric acid. The deep red solution was stirred in the ice bath for 0.5 hours and at room temperature for an additional 3 hours. The mixture was poured into 300 mL water and extracted with ether (2 x 300 mL). The combined ether extracts were washed twice with saturated aqueous NaHCO_3 , twice with water, and once with brine. The organic layer was dried over MgSO_4 , filtered, and concentrated *in vacuo*. The residue purified by column chromatography on silica gel (35% ethyl acetate / hexanes) to give 1.1 g (18%) of methyl ester intermediate as a white semisolid.

Step 2: 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)propionic acid methyl ester


To a stirred solution in DMF (50 mL) of 2,2-dimethyl-3,3-bis(4-hydroxyphenyl)propionic acid methyl ester (1.1 g, 3.7 mmol), prepared as in step 1, was added Cs_2CO_3 (2.5 g, 7.7 mmol) and the mixture stirred 0.5 hours at room temperature. 2-Chloromethylquinoline (1.37 g, 7.7 mmol) was added as a solid in small portions. The reaction mixture was stirred overnight at room temperature. The mixture was poured into 300 mL of water and extracted with ether (2 X 200 mL). Brine was added to the aqueous layer and it was extracted twice with ether. The combined ether extracts were washed with brine, dried over MgSO_4 , filtered, and concentrated *in vacuo* to give a light yellow oil. Chromatography on silica gel (40% ethyl acetate / hexanes) gave 1.68 g (78%) of 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)propionic acid methyl ester as a white foam.

Step 3: 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)propionic acid

- To a solution in methanol (8 mL) and THF (4 mL) of 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)propionic acid methyl ester (0.4 g, 0.69 mmol), prepared as in step 2, was added LiOH·H₂O (4 mL, 4 mmole, 1 M solution) dropwise, and the mixture was stirred overnight after which an additional 2 mL of the LiOH solution was added and the mixture was stirred an additional 24 hours. The mixture was concentrated to dryness, acidified with excess 0.5 M citric acid, diluted with water and extracted with ethyl acetate (3 x 50 mL). The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated *in vacuo*.
- Purification by chromatography on silica gel (7% MeOH/CH₂Cl₂) provided 0.16 g (40%) of 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)propionic acid as a white solid: mp 124-127°C. Anal. Calc'd. for C₃₇H₃₂N₂O₄: C, 78.14; H, 5.67; N 4.92. Found C, 77.45; H, 5.74; N, 4.67.

Step 4: 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)propionic acid sodium salt

- To a solution in THF (10 mL) and ethanol (8 mL) of 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)propionic acid (0.128 g, .225 mmol), prepared as in step 3, was added one equivalent of NaOH (2.3 mL, 0.1 N NaOH). The reaction was stirred for 1 hour at room temperature, concentrated, and dried under high vacuum to provide 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)propionic acid sodium salt as a white powder 0.13 g (99%): mp 250-255°C (dec); ¹H NMR (300 MHz, DMSO-d₆) δ 0.91 (s, 6H), 4.46 (s, 1H), 5.29 (s, 4H), 6.87 (d, 4H, J = 9 Hz), 7.15 (d, 4H, J = 9 Hz), 7.61 (m, 2H), 7.67 (d, 4H, J = 9 Hz), 7.78 (m, 2H), 8.00 (t, 4H, J = 9 Hz), 8.40 (d, 2H, J = 9 Hz); MS m/e (FAB+) 569 (M+H)⁺, (FAB-) 567 (M-H)⁻. Anal. Calc'd. for C₃₇H₃₁N₂O₄Na·0.5 H₂O: C, 74.10; H, 5.37; N 4.67. Found C, 74.02; H, 5.24; N, 4.50.

Example 27**Preparation of [2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)prop-1-yl]oximinoacetic acid sodium salt**

Step 1: 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)propan-1-ol

To a stirred solution in THF (50 mL) at ambient temperature of 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)propionic acid methyl ester (1.33 g, 2.28 mmole), prepared as in Example 26, step 2, was added LiAlH₄ (0.09 g, 2.5 mmol) in a single portion. The mixture was stirred 3 hours at ambient temperature. Water (0.1 mL) was added followed by aqueous 1N NaOH (0.1 ml) and water (0.5 mL). The mixture was then concentrated to dryness and partitioned between water and ethyl acetate. The organic layer was washed with brine, dried over MgSO₄, filtered, and concentrated *in vacuo*. Purification by chromatography on silica gel (60% ethylacetate / hexanes) gave 1.01 g (80%) of 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)-propan-1-ol as a yellow foam.

Step 2: 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)propanal

To a -78 °C solution in methylene chloride (15 mL) of 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)propan-1-ol (0.4 g, 0.72 mmol), prepared as in step 1, and DMSO (0.17 g, 2.17 mmol) was added oxalyl chloride (0.14 g, 1.08 mmol) and the reaction mixture was stirred for 0.5 hours. Triethylamine (0.37 g, 3.6 mmol) was added via syringe, the ice bath was removed, and the reaction was allowed to warm to room temperature. The mixture was concentrated *in vacuo* and triturated with dry THF. The THF solution was filtered and washed with additional THF. The THF solution and washings were combined and concentrated *in vacuo* to give crude 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)propanal which was used without further purification for oxime formation.

Step 3: [2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)prop-1-yl]oximinoacetic acid

The desired compound was prepared according to the method of Example 7, step 2, except substituting 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)-propanal, prepared as in step 2, for 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanal.


Step 4: [2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)prop-1-yl]oximinoacetic acid sodium salt

The desired compound was prepared according to the method of Example 2, except substituting [2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)prop-1-yl]oximinoacetic acid, prepared as in step 2, for 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanoic acid: ¹H NMR (300 MHz, DMSO-d₆) δ 1.03 (s, 6H), 3.89 (s, 1H), 4.06 (s, 2H), 5.32 (s, 4H), 6.97 (d, 4H, J = 9 Hz), 7.31 (d, 4H, J = 9 Hz),

7.46 (s, 1H), 7.65 (m, 4H), 7.78 (m, 2H), 8.01 (t, 4H, $J = 9$ Hz), 8.40 (d, 2H, $J = 9$ Hz); MS (FAB+) m/e 648 ($M+Na$)⁺, 626 ($M+H$)⁺.

Example 28


Preparation of 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)-1-propyliminoxacetic acid




The desired compound was prepared according to the method of Example 5, except substituting 2,2-dimethyl-3,3-bis(4-(2-quinolylmethoxy)phenyl)propan-1-ol prepared as in Example 27, step 1, for 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentan-1-ol: Amorphous white solid, softens at 125 °C (decompose over next 40 °C range); ¹H NMR (300 MHz, DMSO-d₆) δ 0.94 (s, 6H), 3.68 (s, 2H), 3.90 (s, 1H), 5.41 (s, 4H), 6.96 (d, 4H, $J = 9$ Hz), 7.36 (d, 4H, $J = 9$ Hz), 7.41 (m, 2H), 7.63 (m, 4H), 7.78 (m, 2H), 8.00 (t, 4H, $J = 9$ Hz), 8.39 (d, 2H, $J = 9$ Hz). Anal. Calc'd. for C₃₉H₃₅N₃O₅N•2.0 H₂O: C, 70.78; H, 5.94; N 6.34. Found C, 71.25; H, 5.28; N 6.39.

Example 29

Preparation of 2,2-bis(4-(2-quinolylmethoxy)phenyl)acetic acid



The title compound was prepared according to the procedure of Example 12, except substituting glyoxylic acid for pyruvic acid: mp 119-128°C; ¹H NMR (300 MHz, DMSO-d₆) δ 4.91 (s, 1H), 5.34 (s, 4H), 7.01 (d, 4H, $J = 7.5$ Hz), 7.23 (d, 4H, $J = 7.5$ Hz), 7.62 (t, 2H, $J = 7.5$ Hz), 7.66 (d, 2H, $J = 7.5$ Hz), 7.79 (t, 2H, $J = 7.5$ Hz), 8.01 (t, 4H, $J = 7.5$ Hz), 8.40 (d, 2H, $J = 7.5$ Hz). MS (DCI-NH₃) m/e 527 ($M+H$)⁺. Anal. Calc'd. for C₃₄H₂₆N₂O₄•H₂O: C, 74.98; H, 5.18; N, 5.14. Found: C, 74.97; H, 4.75; N, 5.02.

Example 30**Preparation of 2,2-bis(4-(6-fluoro-2-quinolymethoxy)phenyl)acetic acid**


5

The title compound was prepared according to the procedure of Example 12, except substituting glyoxylic acid for pyruvic acid, and substituting 2-chloromethyl-6-fluoroquinoline for 2-chloromethylquinoline: mp 221-223°C; ¹H NMR (300 MHz, DMSO-d₆) δ 4.92 (s, 1H), 5.33 (s, 4H), 7.01 (d, 4H, J = 7.5 Hz), 7.23 (d, 4H, J = 7.5 Hz), 7.70 (m, 4H), 7.82 (dd, 2H, J = 2.5, 9.0 Hz), 8.08 (m, 2H), 8.40 (d, 2H, J = 7.5 Hz); MS (DCI-NH₃) m/e 563(M+H)⁺. Anal. Calc'd. for C₃₄H₂₆N₂O₄: C, 72.59; H, 4.30; N, 4.97. Found: C, 72.36; H, 4.22; N, 4.76.

10

Example 31**Preparation of 2,2-bis(4-(2-quinolymethoxy)phenyl)eth-1-yloximinoacetic acid**

15

**Step 1: 2,2-bis(4-(2-quinolymethoxy)phenyl)acetic acid methyl ester**

20

The desired compound was prepared according to the method of Example 1, steps 1 and 2, except substituting 2,2-bis(4-hydroxyphenyl)acetic acid, prepared as in Example 29, for 4,4-bis(4-hydroxyphenyl)pentanoic acid.

Step 2: 2,2-bis(4-(2-quinolymethoxy)phenyl)eth-1-yloximinoacetic acid

25

The title compound was prepared according to the procedure of Example 27, steps 1-3, except substituting 2,2-bis(4-hydroxyphenyl)acetic acid methyl ester, prepared as in step 1, for methyl 2,2-dimethyl-3,3-bis(4-(2-quinolymethoxy)phenyl)-propionate: mp 104-108°C; ¹H NMR (300 MHz, DMSO-d₆) δ 4.18 (s, 2H), 4.72 (d, 1H, J = 7.5 Hz), 5.33 (s, 4H), 7.01 (d, 4H, J = 7.5 Hz), 7.28 (d, 4H, J = 7.5 Hz), 7.64 (m, 4H), 7.78 (t, 2H, J = 7.5 Hz), 7.98 (m, 5H), 8.40 (d, 2H, J = 7.5 Hz); MS (FAB+) m/e 584 (M+H)⁺, (FAB-) m/e 582 (M-

30

H)⁻. Anal. Calc'd. for C₃₆H₂₉N₃O₅•1.5 H₂O: C, 70.80; H, 5.28; N, 6.88. Found: C, 71.07; H, 5.03; N, 6.61.

Example 32


5 Preparation of 2,2-bis(4-(2-quinolylmethoxy)-phenyl)eth-1-yloximinoacetic acid sodium salt

The desired compound was prepared according to the method of Example 2, except substituting 2,2-bis(4-(2-quinolylmethoxy)phenyl)-1-ethyloximinoacetic acid, prepared as in Example 31, for 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanoic acid: 10 mp 138-145°C; ¹H NMR (300 MHz, DMSO-d₆) δ 4.04 (s, 2H), 4.12 (s, 1H), 4.72 (d, 1H, J = 7.5 Hz), 5.33 (s, 4H), 7.01 (d, 4H, J = 7.5 Hz), 7.28 (d, 4H, J = 7.5 Hz), 7.64 (m, 4H), 7.78 (t, 2H, J = 7.5 Hz), 8.00 (m, 4H), 8.40 (d, 2H, J = 7.5 Hz). MS (DCI-NH₃) m/e 606 (M+Na)⁺, 584 (M+H)⁺.

15

Example 33

Preparation of 3,3-bis(4-(2-quinolylmethoxy)phenyl)propionic acid



20 The desired compound was prepared according to the procedure of Example 26, except substituting ethyl 3,3-diethoxypropionate in ethanol for methyl 3-carboxaldehyde-2,2-dimethyl propionate in methanol: mp 94-111°C; ¹H NMR (300 MHz, DMSO-d₆) δ 2.93 (d, 2H, J = 7.5 Hz), 4.31 (t, 1H, J = 7.5 Hz), 5.31 (s, 4H), 6.95 (d, 4H, J = 8.0 Hz), 7.23 (d, 4H, J = 8.0 Hz), 7.62 (m, 4H), 7.78 (t, 2H, J = 7.5 Hz), 8.00 (t, 4H, J = 7.5 Hz), 8.40 (d, 2H, J = 7.5 Hz); MS (DCI-NH₃) m/e 541 (M+H)⁺. Anal. Calc'd. for C₃₅H₂₈N₂O₄•H₂O: C, 75.25; H, 5.41; N, 5.01. Found: C, 75.09; H, 5.19; N, 4.93.

Example 34

30 Preparation of 3,3-bis(4-(2-quinolylmethoxy)phenyl)propionic acid sodium salt


The desired compound was prepared according to the method of Example 2, except substituting 3,3-bis(4-(2-quinolylmethoxy)phenyl)propionic acid, prepared as in Example 33, for 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanoic acid: mp 224-232°C; ¹H NMR (300 MHz, DMSO-d₆) δ 2.44 (d, 2H, J = 7.5 Hz), 4.31 (t, 1H, J =

7.5 Hz), 5.29 (s, 4H), 6.89 (d, 4H, $J = 9.0$ Hz), 7.12 (d, 4H, $J = 9.0$ Hz), 7.62 (m, 4H), 7.78 (t, 2H, $J = 7.5$ Hz), 8.00 (t, 4H, $J = 7.5$ Hz), 8.40 (d, 2H, $J = 7.5$ Hz); MS (DCI-NH₃) m/e 541 (M+H)⁺.

5

Example 35

Preparation of 4,4-bis(4-(2-quinolylmethoxy)-phenyl)acetic acid-*N*-carboxymethyl amide



10

Step 1: 2,2-bis(4-hydroxyphenyl)acetic acid-*N*-carboxymethyl amide


To a stirred solution in CH₂Cl₂ (0.5 mL), THF (20 mL), and pyridine (20 mL) of 2,2-bis(4-hydroxyphenyl)acetic acid (0.94 g, 3.85 mmol), prepared as in Example 31, was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (1.10 g, 5.76 mmol), glycine methyl ester hydrochloride (0.723 g, 5.76 mmol), and *N*-methylmorpholine (0.63 mL, 5.76 mmol) and the mixture was stirred overnight at room temperature. The reaction mixture was then diluted with ethyl acetate and aqueous 0.5 N HCl. The organic layer was washed with 0.5 N HCl (3X), saturated aqueous NaHCO₃ (3X), and brine, dried over MgSO₄, filtered, and concentrated *in vacuo*. The residue was dried to constant weight on high vacuum to give 1.05 g (87%) of 2,2-bis(4-hydroxyphenyl)acetic acid-*N*-carboxymethyl amide.

20

Step 2: 4,4-bis(4-(2-quinolylmethoxy)phenyl)acetic acid-*N*-carboxymethyl amide

The desired compound was prepared according to the method of Example 1, steps 1-3, except substituting 2,2-bis(4-hydroxyphenyl)acetic acid-*N*-carboxymethyl amide, prepared as in step 1, for 4,4-bis(4-hydroxyphenyl)pentanoic acid: mp 228-231°C; ¹H NMR (300 MHz, DMSO-d₆) δ 3.79 (d, 2H, $J = 6.5$ Hz), 4.93 (s, 1H), 5.34 (s, 4H), 7.00 (d, 4H, $J = 7.5$ Hz), 7.22 (d, 4H, $J = 7.5$), 7.63 (m, 4H), 7.79 (t, 2H, $J = 7.5$ Hz), 8.01 (m, 4H), 8.44 (m, 3H), 12.52 (s, 1H); MS (DCI-NH₃) m/e 584 (M+H)⁺. Anal. Calc'd. for C₃₆H₂₉N₃O₅: C, 74.08; H, 5.01; N, 7.20. Found: C, 73.81; H, 5.20; N, 6.90.

30

Example 36**Preparation of [3,3-bis-(2-quinolylmethoxyphenyl)-
but-1-yl]-2-iminoxypropionic acid**

5

Step 1: 3,3-bis(4-(2-quinolylmethoxy)phenyl)butan-1-ol


To a 0 °C solution in THF (40 mL) of 3,3-bis(4-(2-quinolylmethoxy)phenyl)-
butanoic acid ethyl ester (3.2 g, 5.5 mmol), prepared as in Example 13, was added
10 slowly a solution of LiAlH₄ (1.0 M, 6 mL, 6.0 mmol). The mixture was stirred for 2
hours and water (15 mL) was added slowly followed by aqueous 1N NaOH. The
reaction mixture was diluted with THF (20 mL) and the mixture was filtered through
a celite pad. The filtrate was extracted with ether and the organic extract was washed
with water and brine, dried over MgSO₄, filtered, and concentrated *in vacuo* to
15 provide 3,3-bis(4-(2-quinolylmethoxy)phenyl)butan-1-ol (2.4 g) as a pale yellow
foam.

Step 2: O-[4,4-bis(4-(2-quinolylmethoxy)phenyl)but-1-yl]hydroxylamine

The desired compound was prepared according to the method of Example 5,
20 steps 1 and 2, except substituting 3,3-bis(4-(2-quinolylmethoxy)phenyl)butan-1-ol,
prepared as in step 1, for 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentan-1-ol.

Step 3: [3,3-bis-(2-quinolylmethoxyphenyl)but-1-yl]-2-iminoxypropionic acid

The desired compound was prepared according to the method of Example 6,
25 except substituting O-[4,4-bis(4-(2-quinolylmethoxy)phenyl)but-1-yl]hydroxylamine
for O-[4,4-bis(4-(2-quinolylmethoxy)phenyl)pent-1-yl]hydroxylamine: mp 94-96 °C;
¹H NMR (300 MHz, DMSO-d₆), δ 1.58 (s, 3H), 1.80 (s, 3H), 2.39 (m, 2H), 3.85
(m, 2H), 5.32 (s, 4H), 6.97 (d, 4H, J = 9 Hz), 7.11 (d, 4H, J = 9 Hz), 7.64 (m,
4H), 7.78 (m, 2H), 7.99 (t, 4H, J = 9 Hz), 8.41 (d, 2H, J = 9 Hz). MS (FAB) m/e
30 626 (M+H)⁺. Anal. Calc'd. for: C₃₉H₃₅N₃O₅•H₂O: C, 72.71; H, 5.47; N, 6.53.
Found: C, 72.35; H, 5.42; N, 6.40.

Example 37**Preparation of 4,4-bis(4-(2-quinolylmethoxy)phenyl)-4-hydroxy-2-butynoic acid**

5

Step 1: 4,4-bis(2-quinolylmethoxy)phenyl)-4-hydroxy-2-butynoic acid methyl ester


To a -78 °C solution in THF (40 mL) of *bis*(4-(2-quinolylmethoxy)phenyl) ketone (980 mg, 2 mmol), prepared as in Example 25, step 1, and propiolic acid (0.19 ml, 3 mmol) was added LDA (1.5 M solution in THF, 4 mL, 6 mmol) and the mixture was left at room temperature for 24 hours. The reaction mixture was diluted with water, acidified to pH 5, and extracted with ethyl acetate. The organic phase was dried over MgSO₄, filtered, and concentrated *in vacuo*. The residue was dissolved in DMF (40 mL) and treated with methyl iodide (2 mL) and sodium bicarbonate (170 mg, 2 mmol). The reaction mixture was stirred for 24 hours and then was poured into water. The layers were separated and the aqueous phase was extracted with ethyl acetate. The combined organic phases were washed with brine, dried over MgSO₄, filtered, and concentrated *in vacuo*. Chromatography on silica gel (methylene chloride-ethyl acetate 4:1) provided 730 mg (63 %) of 4,4-*bis*(2-quinolylmethoxy)phenyl)-4-hydroxy-2-butynoic acid methyl ester.

Step 2: 4,4-bis(4-(2-quinolylmethoxy)phenyl)-4-hydroxy-2-butynoic acid

The desired compound was prepared according to the method of Example 1, step 3, except substituting 4,4-*bis*(2-quinolylmethoxy)phenyl)-4-hydroxy-2-butynoic acid methyl ester, prepared as in step 1, for 4,4-*bis*(4-(2-quinolylmethoxy)phenyl)-pentanoic acid: mp 169-172°C; ¹H NMR (300 MHz, DMSO-d₆) δ 5.35 (s, 4H), 7.04 (d, 4H, J = 9 Hz), 7.37 (d, 4H, J = 9 Hz), 7.63 (m, 4H), 7.78 (m, 2H), 8.00 (m, 4H), 8.40 (d, 2H, J = 8 Hz), 13.75 (bs, 1H); MS (DCI-NH₃) m/e: 567 (M+H)⁺. Anal. Calc'd. for C₃₆H₂₆N₂O₄•0.5 H₂O: C, 75.12; H, 4.73; N, 4.89. Found: 75.19; H, 4.80; N, 4.60.

Example 38

Preparation of [5,5-bis(4-(2-quinolylmethoxy)phenyl)-5-hydroxy-3-pentyn-1-yl]-2-iminoxypropionic acid sodium salt



5

Step 1: 5,5-bis(4-(2-quinolylmethoxy)phenyl)-5-hydroxy-3-butyn-1-ol

To a -50 °C solution in THF (25 mL) of *bis*(4-(2-quinolylmethoxy)phenyl) ketone (496 mg, 1 mmol), prepared as in Example 25 was added a solution of dilithium salt of 3-butyn-1-ol (prepared by addition of 1.5 M LDA (3 mL) to 3-butyn-1-ol (0.15 mL, 2 mmol) at -50 °C) and the reaction mixture was allowed to warm to room temperature. The mixture was stirred for 12 hours at room temperature and then was quenched with saturated aqueous ammonium chloride. The layers were separated and the aqueous phase was extracted with ethyl acetate. The combined organic phases were washed with brine, dried over MgSO₄, filtered, and concentrated *in vacuo*. Chromatography on silica gel (silica gel, ethyl acetate) provided 400 mg of 5,5-bis(4-(2-quinolylmethoxy)phenyl)-5-hydroxy-3-butyn-1-ol.

Step 2: [5,5-bis(4-(2-quinolylmethoxy)phenyl)-5-hydroxy-3-pentyn-1-yl]-2-iminoxypropionic acid

The desired compound was prepared according to the method of Example 36, steps 2, and 3, except substituting 5,5-bis(4-(2-quinolylmethoxy)phenyl)-5-hydroxy-3-butyn-1-ol, prepared as in step 1, for 3,3-bis(4-(2-quinolylmethoxy)phenyl)butan-1-ol.

25

Step 3: [5,5-bis(4-(2-quinolylmethoxy)phenyl)-5-hydroxy-3-pentyn-1-yl]-2-iminoxypropionic acid sodium salt

The desired compound was prepared according to the method of Example 2, except substituting [5,5-bis(4-(2-quinolylmethoxy)phenyl)-5-hydroxy-3-pentyn-1-yl]-2-iminoxypropionic acid, prepared as in Step 2, for 4,4-bis(4-(2-quinolylmethoxy)phenyl)pentanoic acid: mp 108-111 °C; ¹H NMR (300 MHz, DMSO-d₆) δ: 1.83 (s, 3H), 2.61 (t, 2H, J = 7 Hz), 4.10 (t, 2H, J = 7 Hz), 5.32 (s, 4H), 6.96 (d, 4H, J = 9 Hz), 7.40 (d, 4H, J = 9 Hz), 7.62 (m, 4H), 7.78 (m, 2H), 8.00 (m, 4H), 8.40 (d, 2H, J = 8 Hz); MS (FAB+) m/e 674 (M+Na)⁺, 652


30

(M+H)⁺, (FAB-) m/e 651 (M-H)⁻. Anal. Calc'd. for C₄₀H₃₂N₃O₆Na•H₂O: C, 69.46; H, 4.95; N, 6.07. Found: C, 69.62; H, 5.10; N, 5.61.

The following additional examples are prepared according to the method described in Example 1, except substituting the requisite heteroarylmethylhalide W-CH₂X where X is Cl, Br, or I for 2-chloromethylquinoline hydrochloride.


Example 39

4,4-bis(4-(2-benzoxazolylmethoxy)phenyl)pentanoic acid




Example 40

4,4-bis(4-(2-pyrimidylmethoxy)phenyl)pentanoic acid




Example 41

4,4-bis(4-(4-phenyl-2-thiazolylmethoxy)phenyl)pentanoic acid



Example 42


4,4-bis(4-(4-(pyrid-2-yl)-2-thiazolylmethoxy)phenyl)pentanoic acid



5

Example 43


4,4-bis(4-(6-phenyl-2-pyridylmethoxy)phenyl)pentanoic acid



10

Example 44


4,4-bis(4-(5-phenyl-2-pyridylmethoxy)phenyl)pentanoic acid



15


Example 45

4,4-bis(4-(6-(pyrid-2-yl))-2-pyridylmethoxy)phenyl)pentanoic acid




Example 46

4,4-bis(4-(4-phenyl-2-pyrimidinylmethoxy)phenyl)pentanoic acid



WHAT IS CLAIMED IS:

1. A compound of formula:



- 5 or a pharmaceutically acceptable salt thereof:

wherein **W** is the same at each occurrence and is selected from the group consisting of

- (a) quinolyl;
 (b) quinolyl substituted with a substituent selected from the group
 10 consisting of:

halogen,
 alkyl of one to six carbon atoms,
 phenyl,
 phenyl substituted with a substituent selected from the group

- 15 consisting of:

halogen,
 alkyl of one to six carbon atoms,
 haloalkyl of one to six carbon atoms, and
 alkoxy of one to six carbon atoms;

20

pyridyl; and

pyridyl substituted with a substituent selected from the group

consisting of:

halogen,
 alkyl of one to six carbon atoms, and,
 alkoxy of one to six carbon atoms,

25

- (c) benzothiazolyl;

- (d) benzothiazolyl substituted with a substituent selected from the group
 consisting of:

halogen,

30

alkyl of one to six carbon atoms,

phenyl.

phenyl substituted with a substituent selected from the group

consisting of:

35 halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,
pyridyl, and
pyridyl substituted with a substituent selected from the group

40 consisting of:

halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;

(e) benzoxazolyl;

45 (f) benzoxazolyl substituted with a substituent selected from the group consisting of:

halogen,
alkyl of one to six carbon atoms,
phenyl,
phenyl substituted with a substituent selected from the group

consisting of:

55 halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,
pyridyl, and
pyridyl substituted with a substituent selected from the group

consisting of:

60 halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;

(g) benzimidazolyl;

(h) benzimidazolyl substituted with a substituent selected from the group

consisting of:

65 halogen,
alkyl of one to six carbon atoms,
phenyl.
phenyl substituted with a substituent selected from the group

consisting of:

- 70 halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,
pyridyl, and
- 75 pyridyl substituted with a substituent selected from the group
consisting of:
- halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;
- 80 (i) quinoxalyl;
(j) quinoxalyl substituted with a substituent selected from the group
consisting of:
- halogen,
alkyl of one to six carbon atoms,
85 phenyl,
phenyl substituted with a substituent selected from the group
consisting of:
- halogen,
alkyl of one to six carbon atoms,
90 haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,
pyridyl, and
pyridyl substituted with a substituent selected from the group
consisting of:
- 95 halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;
- (k) pyridyl;
(l) pyridyl substituted with a substituent selected from the group
- 100 consisting of:
- phenyl,
phenyl substituted with a substituent selected from the group
consisting of:
- 105 halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and

- alkoxy of one to six carbon atoms,
pyridyl, and
pyridyl substituted with a substituent selected from the group
- 110 consisting of:
- halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;
- (m) pyrimidyl;
- 115 (n) pyrimidyl substituted with a substituent selected from the group
consisting of:
- phenyl,
phenyl substituted with a substituent selected from the group
- consisting of:
- 120 halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,
pyridyl, and
- 125 pyridyl substituted with a substituent selected from the group
consisting of:
- halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;
- 130 (o) thiazolyl, and
(p) thiazolyl substituted with a substituent selected from the group
consisting of:
- phenyl,
phenyl substituted with a substituent selected from the group
- 135 consisting of:
- halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,
- 140 pyridyl, and
pyridyl substituted with a substituent selected from the group
consisting of:

(c) $-NR^7R^8$ where R^7 and R^8 are independently selected
180 from the group consisting of:

hydrogen,
alkyl of one to six carbon atoms,
hydroxy, and
alkoxy of one to six carbon atoms,

185 or R^7 and R^8 taken together define a five- to eight-
membered ring, with the proviso that R^7 and R^8 may not simultaneously be
hydroxyl,

(d) $-NR^3SO_2R^9$ wherein R^3 is as defined above and R^9 is
alkyl of one to six carbon atoms,

190 (e) $-NH$ -tetrazolyl, and

(f) glycynyl.

2. A compound or pharmaceutically acceptable salt thereof as
defined by Claim 1 wherein Z is selected from the group consisting of

- (a) COM,
(b) $CH=N-O-A-COM$,
5 (c) $CH_2-O-N=A-COM$, and
(d) OH wherein A is alkylene of one to six carbon atoms, and M is -OH.

3. A compound or pharmaceutically acceptable salt thereof as
defined by Claim 2 wherein W is the same at each occurrence and is selected from the
group consisting of

- (a) pyridyl;
5 (b) pyridyl substituted with a substituent selected from the group
consisting of:

phenyl,
phenyl substituted with a substituent selected from the group

consisting of:

10 halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,

pyridyl, and

15 pyridyl substituted with a substituent selected from the group
consisting of:

- halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;
- 20 (c) pyrimidyl;
(d) pyrimidyl substituted with a substituent selected from the group
consisting of:
- phenyl,
phenyl substituted with a substituent selected from the group
- 25 consisting of:
- halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,
- 30 pyridyl, and
pyridyl substituted with a substituent selected from the group
consisting of:
- halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;
- 35 (e) thiazolyl, and
(f) thiazolyl substituted with a substituent selected from the group
consisting of:
- phenyl,
phenyl substituted with a substituent selected from the group
- 40 consisting of:
- halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,
- 45 pyridyl, and
pyridyl substituted with a substituent selected from the group
consisting of:
- 50 halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms.

4. A compound or pharmaceutically acceptable salt thereof as defined by Claim 2 wherein W is the same at each occurrence and is selected from the group consisting of:

(a) quinolyl;

5 (b) quinolyl substituted with a substituent selected from the group consisting of:

halogen,

alkyl of one to six carbon atoms,

phenyl,

10 phenyl substituted with a substituent selected from the group

consisting of:

halogen,

alkyl of one to six carbon atoms,

haloalkyl of one to six carbon atoms, and

alkoxy of one to six carbon atoms;

15 pyridyl; and

pyridyl substituted with a substituent selected from the group

consisting of:

halogen,

alkyl of one to six carbon atoms, and,

20 alkoxy of one to six carbon atoms,

(c) benzothiazolyl;

(d) benzothiazolyl substituted with a substituent selected from the group consisting of:

halogen,

25 alkyl of one to six carbon atoms,

phenyl,

phenyl substituted with a substituent selected from the group

consisting of:

halogen,

30 alkyl of one to six carbon atoms,

haloalkyl of one to six carbon atoms, and

alkoxy of one to six carbon atoms,

pyridyl, and

pyridyl substituted with a substituent selected from the group

35 consisting of:

- 56 -

halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;

(e) benzoxazolyl;

40 (f) benzoxazolyl substituted with a substituent selected from the group
consisting of:

halogen,
alkyl of one to six carbon atoms,
phenyl,
45 phenyl substituted with a substituent selected from the group

consisting of:

halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
50 alkoxy of one to six carbon atoms,
pyridyl, and
pyridyl substituted with a substituent selected from the group

consisting of:

halogen,
55 alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;

(g) benzimidazolyl;

(h) benzimidazolyl substituted with a substituent selected from the group
consisting of:

60 halogen,
alkyl of one to six carbon atoms,
phenyl,
phenyl substituted with a substituent selected from the group

consisting of:

65 halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,

pyridyl, and

70 pyridyl substituted with a substituent selected from the group
consisting of:

- 57 -

- halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;
- 75 (i) quinoxalyl; and
(j) quinoxalyl substituted with a substituent selected from the group
consisting of:
- halogen,
alkyl of one to six carbon atoms,
80 phenyl,
phenyl substituted with a substituent selected from the group
consisting of:
- halogen,
alkyl of one to six carbon atoms,
85 haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,
pyridyl, and
pyridyl substituted with a substituent selected from the group
consisting of:
- 90 halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms.

5. A compound or pharmaceutically acceptable salt thereof as defined by Claim 4 wherein W is the same at each occurrence and is selected from the group consisting of

- (a) quinolyl;
- 5 (b) quinolyl substituted with a substituent selected from the group
consisting of:
- halogen,
alkyl of one to six carbon atoms,
phenyl,
10 phenyl substituted with a substituent selected from the group
consisting of:
- halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
15 alkoxy of one to six carbon atoms;

- pyridyl; and
pyridyl substituted with a substituent selected from the group
consisting of:
- 20 halogen,
alkyl of one to six carbon atoms, and,
alkoxy of one to six carbon atoms,
- (c) benzothiazolyl; and
(d) benzothiazolyl substituted with a substituent selected from the group
consisting of:
- 25 halogen,
alkyl of one to six carbon atoms,
phenyl,
phenyl substituted with a substituent selected from the group
consisting of:
- 30 halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,
pyridyl, and
35 pyridyl substituted with a substituent selected from the group
consisting of:
- halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;

6. A compound or pharmaceutically acceptable salt thereof as
defined by Claim 5 wherein W is the same at each occurrence and is selected from the
group consisting of:
- (a) quinolyl; and
5 (b) quinolyl substituted with a substituent selected from the group
consisting of:
- halogen,
alkyl of one to six carbon atoms,
phenyl,
10 phenyl substituted with a substituent selected from the group
consisting of:

halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;
pyridyl; and
pyridyl substituted with a substituent selected from the group

consisting of:

20 halogen,
alkyl of one to six carbon atoms, and,
alkoxy of one to six carbon atoms.

7. A compound or pharmaceutically acceptable salt thereof as defined by Claim 1 wherein Z is COM wherein M is -OH.

8. A compound or pharmaceutically acceptable salt thereof as defined by Claim 7 wherein X is alkylene of one to six carbon atoms.

9. A compound or pharmaceutically acceptable salt thereof as defined by Claim 8 wherein W is the same at each occurrence and is selected from the group consisting of:

(a) pyridyl;

5 (b) pyridyl substituted with a substituent selected from the group consisting of:

phenyl.
phenyl substituted with a substituent selected from the group
consisting of:

10 halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,
pyridyl, and
15 pyridyl substituted with a substituent selected from the group
consisting of:

20 (c) pyrimidyl;
halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;

- 60 -

(d) pyrimidyl substituted with a substituent selected from the group consisting of:

phenyl,
phenyl substituted with a substituent selected from the group
25 consisting of:

halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,
30 pyridyl, and
pyridyl substituted with a substituent selected from the group
consisting of:

halogen,
alkyl of one to six carbon atoms, and
35 alkoxy of one to six carbon atoms;

(e) thiazolyl, and

(f) thiazolyl substituted with a substituent selected from the group
consisting of:

phenyl,
phenyl substituted with a substituent selected from the group
40 consisting of:

halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
45 alkoxy of one to six carbon atoms,
pyridyl, and
pyridyl substituted with a substituent selected from the group
consisting of:

halogen,
50 alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms.

10. A compound or pharmaceutically acceptable salt thereof as defined by Claim 8 wherein W is the same at each occurrence and is selected from the group consisting of:

(a) quinolyl;

5 (b) quinolyl substituted with a substituent selected from the group
consisting of:

 halogen,
 alkyl of one to six carbon atoms,
 phenyl,
10 phenyl substituted with a substituent selected from the group

consisting of:

 halogen,
 alkyl of one to six carbon atoms,
 haloalkyl of one to six carbon atoms, and
15 alkoxy of one to six carbon atoms;
 pyridyl; and
 pyridyl substituted with a substituent selected from the group

consisting of:

 halogen,
20 alkyl of one to six carbon atoms, and,
 alkoxy of one to six carbon atoms,

(c) benzothiazolyl;

(d) benzothiazolyl substituted with a substituent selected from the group
consisting of:

25 halogen,
 alkyl of one to six carbon atoms,
 phenyl,
 phenyl substituted with a substituent selected from the group

consisting of:

30 halogen,
 alkyl of one to six carbon atoms,
 haloalkyl of one to six carbon atoms, and
 alkoxy of one to six carbon atoms,
 pyridyl, and
35 pyridyl substituted with a substituent selected from the group

consisting of:

 halogen,
 alkyl of one to six carbon atoms, and
 alkoxy of one to six carbon atoms;

40 (e) benzoxazolyl;

(f) benzoxazolyl substituted with a substituent selected from the group consisting of:

45 halogen,
alkyl of one to six carbon atoms,
phenyl,
phenyl substituted with a substituent selected from the group

consisting of:

50 halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,
pyridyl, and
pyridyl substituted with a substituent selected from the group

consisting of:

55 halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;

(g) benzimidazolyl;

60 (h) benzimidazolyl substituted with a substituent selected from the group consisting of:

halogen,
alkyl of one to six carbon atoms,
phenyl,
phenyl substituted with a substituent selected from the group

65 consisting of:

halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,
70 pyridyl, and
pyridyl substituted with a substituent selected from the group

consisting of:

75 halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;

(i) quinoxalyl; and

(j) quinoxalyl substituted with a substituent selected from the group consisting of:

80 halogen,
 alkyl of one to six carbon atoms,
 phenyl,
 phenyl substituted with a substituent selected from the group

consisting of:

85 halogen,
 alkyl of one to six carbon atoms,
 haloalkyl of one to six carbon atoms, and
 alkoxy of one to six carbon atoms,
 pyridyl, and
 pyridyl substituted with a substituent selected from the group

90 consisting of:

 halogen,
 alkyl of one to six carbon atoms, and
 alkoxy of one to six carbon atoms

11. A compound or pharmaceutically acceptable salt thereof as defined by Claim 8 wherein W is the same at each occurrence and is selected from the group consisting of:

5 (a) quinolyl;
 (b) quinolyl substituted with a substituent selected from the group consisting of:

 halogen,
 alkyl of one to six carbon atoms,
 phenyl,
10 phenyl substituted with a substituent selected from the group

consisting of:

 halogen,
 alkyl of one to six carbon atoms,
 haloalkyl of one to six carbon atoms, and
15 alkoxy of one to six carbon atoms;
 pyridyl; and
 pyridyl substituted with a substituent selected from the group

consisting of:

- 64 -

- 20 halogen,
alkyl of one to six carbon atoms, and,
alkoxy of one to six carbon atoms,
- (c) benzothiazolyl; and
- (d) benzothiazolyl substituted with a substituent selected from the group
consisting of:
- 25 halogen,
alkyl of one to six carbon atoms,
phenyl,
phenyl substituted with a substituent selected from the group
consisting of:
- 30 halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms,
pyridyl, and
- 35 pyridyl substituted with a substituent selected from the group
consisting of:
- 40 halogen,
alkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms.
12. A compound or pharmaceutically acceptable salt thereof as
defined by Claim 11 wherein W is the same at each occurrence and is selected from
the group consisting of:
- (a) quinolyl; and
- 5 (b) quinolyl substituted with a substituent selected from the group
consisting of:
- 10 halogen,
alkyl of one to six carbon atoms,
phenyl,
phenyl substituted with a substituent selected from the group
consisting of:
- 15 halogen,
alkyl of one to six carbon atoms,
haloalkyl of one to six carbon atoms, and
alkoxy of one to six carbon atoms;

pyridyl; and

pyridyl substituted with a substituent selected from the group

consisting of:

halogen,

20

alkyl of one to six carbon atoms, and,

alkoxy of one to six carbon atoms.

13. A compound or pharmaceutically acceptable salt thereof
selected from the group consisting of:

4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid,

4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid sodium salt,

5 4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid magnesium salt,

4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentan-1-ol,

[4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pent-1-yl]iminoxyacetic acid,

[4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pent-1-yl]-2-iminoxypropionic acid,

[4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pent-1-yl]oximinoacetic acid,

10 4,4-*bis*(4-(2-benzothiazolylmethoxy)phenyl)pentanoic acid,

4,4-*bis*(4-(2-benzothiazolylmethoxy)phenyl)pentanoic acid sodium salt,

4,4-*bis*(4-(7-chloro-2-quinolylmethoxy)phenyl)pentanoic acid,

4,4-*bis*(4-(7-fluoro-2-quinolylmethoxy)phenyl)pentanoic acid sodium salt,

2,2-*bis*(4-(2-quinolylmethoxy)phenyl)propionic acid,

15 3,3-*bis*(4-(2-quinolylmethoxy)phenyl)butanoic acid,

5,5-*bis*(4-(2-quinolylmethoxy)phenyl)hexanoic acid,

5,5-*bis*(4-(2-quinolylmethoxy)phenyl)hexanoic acid sodium salt,

4,4-*bis*-(4-(2-pyridylmethoxy)phenyl)pentanoic acid sodium salt,

4,4-*bis*-(4-(2-quinolylmethoxy)phenyl)pentanoic acid sodium salt,

20 4,4-*bis*(4-(1-methyl-2-benzimidazolylmethoxy)phenyl)pentanoic acid,

4,4-*bis*(4-(1-methyl-2-benzimidazolylmethoxy)phenyl)pentanoic acid sodium

salt,

4,4-*bis*(3-chloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid,

4,4-*bis*(3-chloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid sodium salt,

25 [4,4-*bis*(3-chloro-4-(2-quinolylmethoxy)phenyl)pent-1-yl]-2-

iminoxypropionic acid sodium salt,

4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid *N,N*-diethylhydroxyl
amine-*O*-ester,

2,2-*bis*(4-(2-quinolylmethoxy)phenyl)butyric acid,

30 1,1-*bis*(4-(2-quinolylmethoxy)phenyl)ethanol,

2,2-dimethyl-3,3-*bis*(4-(2-quinolylmethoxy)phenyl)propionic acid sodium salt,
 [2,2-dimethyl-3,3-*bis*(4-(2-quinolylmethoxy)phenyl)prop-1-yl]oximinoacetic acid sodium salt,
 35 2,2-dimethyl-3,3-*bis*(4-(2-quinolylmethoxy)phenyl)-1-propyliminoxyacetic acid,
 2,2-*bis*(4-(2-quinolylmethoxy)phenyl)acetic acid,
 2,2-*bis*(4-(6-fluoro-2-quinolylmethoxy)phenyl)acetic acid,
 2,2-*bis*(4-(2-quinolylmethoxy)phenyl)eth-1-yloximinoacetic acid,
 40 2,2-*bis*(4-(2-quinolylmethoxy)phenyl)eth-1-yloximinoacetic acid sodium salt,
 3,3-*bis*(4-(2-quinolylmethoxy)phenyl)propionic acid,
 3,3-*bis*(4-(2-quinolylmethoxy)phenyl)propionic acid sodium salt,
 4,4-*bis*(4-(2-quinolylmethoxy)phenyl)acetic acid-*N*-carboxymethyl amide,
 [3,3-*bis*-(2-quinolylmethoxyphenyl)but-1-yl]-2-iminoxypropionic acid,
 45 4,4-*bis*(4-(2-quinolylmethoxy)phenyl)-4-hydroxy-2-butynoic acid, and
 [5,5-*bis*(4-(2-quinolylmethoxy)phenyl)-5-hydroxy-3-pentyn-1-yl]-2-iminoxypropionic acid sodium salt.

14. A compound or pharmaceutically acceptable salt thereof selected from the group consisting of:

4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid,
 4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid sodium salt,
 5 4,4-*bis*(4-(2-quinolylmethoxy)phenyl)pentanoic acid magnesium salt,
 4,4-*bis*(4-(2-benzothiazoylmethoxy)phenyl)pentanoic acid,
 4,4-*bis*(4-(2-benzothiazoylmethoxy)phenyl)pentanoic acid sodium salt,
 4,4-*bis*(4-(7-chloro-2-quinolylmethoxy)phenyl)pentanoic acid,
 4,4-*bis*(4-(7-fluoro-2-quinolylmethoxy)phenyl)pentanoic acid sodium salt,
 10 2,2-*bis*(4-(2-quinolylmethoxy)phenyl)propionic acid,
 3,3-*bis*(4-(2-quinolylmethoxy)phenyl)butanoic acid,
 5,5-*bis*(4-(2-quinolylmethoxy)phenyl)hexanoic acid,
 5,5-*bis*(4-(2-quinolylmethoxy)phenyl)hexanoic acid sodium salt,
 4,4-*bis*(3-chloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid,
 15 4,4-*bis*(3-chloro-4-(2-quinolylmethoxy)phenyl)pentanoic acid sodium salt,
 2,2-*bis*(4-(2-quinolylmethoxy)phenyl)butyric acid,
 2,2-dimethyl-3,3-*bis*(4-(2-quinolylmethoxy)phenyl)propionic acid sodium salt,
 2,2-*bis*(4-(2-quinolylmethoxy)phenyl)acetic acid,

- 20 2,2-*bis*(4-(6-fluoro-2-quinolymethoxy)phenyl)acetic acid,
 2,2-*bis*(4-(2-quinolymethoxy)phenyl)propionic acid, and
 2,2-*bis*(4-(2-quinolymethoxy)phenyl)propionic acid sodium salt.

15. A method for inhibiting lipoxygenase activity or leukotriene biosynthesis in a mammal in need of such treatment comprising administering to the mammal a therapeutically effective amount of a compound of Claim 1.

16. A composition for inhibiting lipoxygenase activity or the biosynthesis of leukotrienes comprising a pharmaceutical carrier and a therapeutically effective amount of a compound of Claim 1.

INTERNATIONAL SEARCH REPORT

Inter. Application No

PCT/US 96/15345

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D215/14 A61K31/395 C07D277/64 C07D215/18 C07D213/30
C07D241/42 C07D235/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94 27968 A (MERCK FROSST CANADA INC.) 8 December 1994 cited in the application see claims ---	1,16
A	EP 0 349 062 A (MERCK FROSST CANADA INC.) 3 January 1990 cited in the application see claims ---	1,16
A	US 4 970 215 A (BAYER AG) 13 November 1990 cited in the application see claims -----	1,16

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

5 December 1996

Date of mailing of the international search report

20.12.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/ 15345

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 15 is directed to a method of treatment of (diagnostic method practised on) the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 96/15345

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9427968	08-12-94	US-A- 5438141	01-08-95
		AU-A- 6791294	20-12-94
		EP-A- 0699188	06-03-96

EP-A-349062	03-01-90	CA-A- 1333714	27-12-94
		JP-A- 2076855	16-03-90
		US-A- 5102881	07-04-92
		US-A- 5232916	03-08-93

US-A-4970215	13-11-90	DE-A- 3900261	07-12-89
		AU-B- 616269	24-10-91
		AU-A- 3527089	07-12-89
		CA-A- 1333802	03-01-95
		CN-A- 1038641	10-01-90
		CN-B- 1030251	15-11-95
		DK-B- 169544	28-11-94
		EP-A- 0344519	06-12-89
		ES-T- 2053864	01-08-94
		FI-B- 91635	15-04-94
		HK-A- 38895	24-03-95
		HU-A- 9500147	28-07-95
		IE-B- 61922	30-11-94
		JP-A- 2019359	23-01-90
		NO-B- 174889	18-04-94
		PT-B- 90675	31-10-94
		SG-A- 12795	16-06-95

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.